

Bifibrations of Polycategories and Classical Linear Logic

Extended version

Nicolas Blanco¹

*Computer Science Department
University of Birmingham
Birmingham, UK*

Noam Zeilberger²

*Laboratoire d'Informatique de l'École Polytechnique
Palaiseau, France*

Abstract

The main goal of this article is to expose and relate different ways of interpreting the multiplicative fragment of classical linear logic in polycategories. Polycategories are known to give rise to models of classical linear logic in so-called representable polycategories with duals, which ask for the existence of various polymaps satisfying the different universal properties needed to define tensor, par, and negation. We begin by explaining how these different universal properties can all be seen as instances of a single notion of universality of a polymap parameterised by an input or output object, which also generalises the classical notion of universal multimap in a multicategory. We then proceed to introduce a definition of in-cartesian and out-cartesian polymaps relative to a refinement system (= strict functor) of polycategories, in such a way that universal polymaps can be understood as a special case. In particular, we obtain that a polycategory is a representable polycategory with duals if and only if it is bifibred over the terminal polycategory $\mathbb{1}$. Finally, we present a Grothendieck correspondence between bifibrations of polycategories and pseudofunctors into \mathbf{MAj} , the (weak) 2-polycategory of multivariable adjunctions. When restricted to bifibrations over $\mathbb{1}$ we get back the correspondence between *-autonomous categories and Frobenius pseudomonoids in \mathbf{MAj} that was recently observed by Shulman.

Keywords: Polycategories, linear logic, bifibrations, Grothendieck construction, Frobenius monoids

1 Introduction

In his early studies of the linguistic applications of Gentzen's sequent calculus [Lam61], Lambek observed that the so-called "associative syntactic calculus" of [Lam58] has a natural semantic interpretation, where formulas are interpreted as bimodules of rings and proofs of sequents $A_1, \dots, A_n \rightarrow B$ are interpreted as multilinear maps $A_1 \times \dots \times A_n \rightarrow B$. He mentions that one benefit of the sequent calculus presentation is that it leads to a decision procedure for the existence of canonical mappings, and notes that "it has already been observed by Bourbaki [*Algèbre multilinéaire*, 1948] that linear mappings of the kind we are interested in are best defined with the help of multilinear mappings". These early observations later led Lambek to formally introduce the definition of *multicategories* in [Lam69], which generalise categories by allowing morphisms to have multiple inputs, a paradigmatic example being the multicategory of vector spaces and multilinear maps.

Szabo, a student of Lambek, introduced *polycategories* in [Sza75], which further generalise multicategories by allowing morphisms to have multiple outputs in addition to multiple inputs. One motivation for studying polycategories from the view of proof theory is that they stand in the same relation to Gentzen's classical

¹ Email: n.blanco@pgr.bham.ac.uk

² Email: noam.zeilberger@lix.polytechnique.edu

sequent calculus LK as multicategories stand in relation to the intuitionistic sequent calculus LJ. For example, the composition operation for morphisms in a polycategory is typed just like the cut rule in classical sequent calculus. Lambek and Szabo’s work was later revisited from the perspective of linear logic [Gir87] by Cockett and Seely [CS97], see also [BCST96,Hyl02,CS07]. In particular, the notion of a *representable (or two-tensor) polycategory with duals* provides a natural source of models for the multiplicative fragment of classical linear logic. Representable polycategories with duals are equivalent to the $*$ -autonomous categories of Barr [Bar91], but have the advantage that all of the logical connectives can be defined by the existence of objects and (poly)morphisms satisfying certain universal properties, rather than as algebraic structures subject to coherence conditions.

This relation between $*$ -autonomous categories and representable polycategories with duals is analogous to the relation between monoidal categories and *representable multicategories* (called *monoidal multicategories* by Lambek [Lam69]), a relation studied carefully by Hermida [Her00]. Hermida noted certain analogies between the theory of representable multicategories and the theory of fibred categories (cf. [Her00, Table 1]), which he later made explicit by introducing a notion of (covariant) *fibration of multicategories* [Her04], in such a way that a representable multicategory is precisely the same thing as a multicategory fibred over the terminal multicategory $\mathbb{1}$. One interest of studying the more general notion of covariant fibration of multicategories $\mathcal{E} \rightarrow \mathcal{B}$, where every multimorphism $f : A_1, \dots, A_n \rightarrow B$ in \mathcal{B} induces a pushforward functor $\mathbf{push}\langle f \rangle : \mathcal{E}_{A_1} \times \dots \times \mathcal{E}_{A_n} \rightarrow \mathcal{E}_B$, is that it models a much richer class of structures coming from algebra and logic. For example, Hermida notes that an *algebra for an operad* O can be identified with a discrete covariant fibration over O , the latter seen as a one-object multicategory. The appropriate definition of *contravariant* fibration (and of bifibration) of multicategories was not addressed in [Her04]. However, there is a natural definition of contravariant fibration of multicategories, made explicit in the work of Hörmann [H17, A.2] and of Licata, Shulman, and Riley [LSR17], under which each multimorphism of the base multicategory induces a family of pullback operations $\mathbf{pull}[f]^{(i)} : \mathcal{E}_{A_1}^{\text{op}} \times \dots \times \mathcal{E}_{A_{i-1}}^{\text{op}} \times \mathcal{E}_{A_{i+1}}^{\text{op}} \times \dots \times \mathcal{E}_{A_n}^{\text{op}} \times \mathcal{E}_B \rightarrow \mathcal{E}_{A_i}$, parameterised by the selection of the index $1 \leq i \leq n$ of a particular input object A_i . One interesting feature of this definition is that *monoidal biclosed categories* in the sense of Lambek [Lam69] are equivalent to multicategories bifibred over $\mathbb{1}$. Moreover, replacing the terminal multicategory by an arbitrary base multicategory leads to a much richer framework for modelling a variety of substructural and modal logics, as discussed by Licata et al. [LSR17], and in a very similar spirit to Melliès and Zeilberger’s work on type refinement and monoidal closed bifibrations (cf. [MZ15,MZ16]). In particular, a recurring pattern is that some algebraic gadget in the base (e.g., a monoid object) induces some logical structure (e.g., monoidal closure) on its fibre.

In this paper, we begin to develop a theory of bifibrations of polycategories, guided by the principle that representable polycategories with duals (and hence $*$ -autonomous categories) should be equivalent to polycategories bifibred over the terminal polycategory $\mathbb{1}$. One consequence of this theory is that we recover a nice observation recently made by Shulman [Shu19], that $*$ -autonomous categories are equivalent to (pseudo) Frobenius monoids in the (2-)polycategory of multivariable adjunctions. This will follow as a result of a general Grothendieck construction for bifibrations of polycategories, in a similar manner to the pattern mentioned above. Perhaps surprisingly, another one of our original motivations for developing this theory was trying to better understand properties of the category \mathbf{FBan}_1 of finite dimensional Banach spaces and contractive maps. It is a $*$ -autonomous category and it comes with a $*$ -autonomous forgetful functor into \mathbf{FVect} , but contrary to the latter it is not compact closed. It provides a model of classical MALL based on finite dimensional vector spaces that is not degenerate, in the sense that the positive and negative fragments do not coincide. While the tensor, and more generally the use of \mathbf{FBan}_1 as a model of intuitionistic MALL is well-documented (cf. [BPS94]) we could not find any mention of the par in the literature. In fact, this category is one of the original examples of $*$ -autonomous category provided by Barr in [Bar79, Ch. 4, 53–59], but without describing the tensor and par in \mathbf{FBan}_1 explicitly. Yet, the structures needed to interpret them are popular in the study of Banach spaces: \otimes and \wp correspond to different norms placed on the tensor product of vector spaces called the *projective* and the *injective* (cross)norms, which have the property of being extremal in all the well-behaved norms that can be put on the tensor product. More specifically for any crossnorm $\| - \|$ and any $u \in A \otimes B$ we have $\|u\|_{A\wp B} \leq \|u\| \leq \|u\|_{A\otimes B}$. We will see that this has a nice explanation from the fact that the projective (\otimes) norm and the injective (\wp) norm can be defined as pushforwards and pullbacks, respectively, relative to the forgetful functor into vector spaces.

2 Polycategories, linear logic, and universality

2.1 Polycategories

There are several different definitions of “polycategory” in the literature. We will consider the following definition of (non-symmetric) polycategory due to Cockett and Seely [CS97], which differs slightly from Szabo’s original definition [Sza75] in imposing a *planarity condition* on composition. The ideas in this paper may be transferred in an almost straightforward way to the setting of symmetric polycategories (cf. [Hyl02,Shu20]),

but we work with planar polycategories for the sake of greater generality.

Definition 2.1 A *polycategory* \mathcal{P} consists of:

- a collection of *objects* $Ob(\mathcal{P})$
- for any pair of finite lists of objects Γ and Δ , a set $\mathcal{P}(\Gamma; \Delta)$ of *polymaps* from Γ to Δ denoted $f: \Gamma \rightarrow \Delta$ (we refer to objects in Γ as *inputs* of f , and to objects in Δ as *outputs*)
- for every object A , an *identity* polymap $id_A: A \rightarrow A$
- for any pair of polymaps $f: \Gamma \rightarrow \Delta_1, A, \Delta_2$ and $g: \Gamma'_1, A, \Gamma'_2 \rightarrow \Delta'$ satisfying the restriction that [either Δ_1 or Γ'_1 is empty] and [either Δ_2 or Γ'_2 is empty], a polymap $g \circ_A f: \Gamma'_1, \Gamma, \Gamma'_2 \rightarrow \Delta_1, \Delta', \Delta_2$

subject to appropriate *unitality*, *associativity*, and *interchange* laws whenever these make sense:

$$id_A \circ_A f = f \tag{1}$$

$$f \circ_A id_A = f \tag{2}$$


$$(h \circ_B g) \circ_A f = h \circ_B (g \circ_A f) \tag{3}$$

$$(h \circ_B g) \circ_A f = (h \circ_A f) \circ_B g \tag{4}$$


$$h \circ_B (g \circ_A f) = g \circ_A (h \circ_B f) \tag{5}$$

Remark 2.2 The notation \circ_A for the composition can be ambiguous when there are multiple copies of the same object. This can be dealt with more carefully by indexing or labelling each input and output of a polymap. However, we will stick with the more relaxed (albeit less precise) notation in this article, since it will never lead to ambiguity in the examples.

Remark 2.3 We will sometimes find it useful to represent polymaps by string diagrams. In this diagrammatic syntax, the composition operation may be depicted schematically as follows:



The restriction on the composition operation that either Δ_1 or Γ'_1 is empty and that either Δ_2 or Γ'_2 is empty is called a “planarity” condition, since in the picture above it means that there are actually no crossing wires. In general, the string diagram of a polymap corresponds to a planar tree with the edges oriented from left to right, and the polycategory axioms correspond to natural isotopies between diagrams. For example, the interchange law (4) states that when composing along two different inputs, the order should not matter:



This justifies drawing the two polymaps f and g above on the same level, as we will sometimes do in examples.

2.2 Representable polycategories with duals

In this section we briefly recall the notion of representable (or two-tensor) polycategory with duals, which has been used to model the multiplicative connectives of classical linear logic.

Definition 2.4 Let Γ be a list of objects in a polycategory \mathcal{P} . A *tensor product* of Γ is an object $\otimes \Gamma$ equipped with a polymap $m_\Gamma: \Gamma \rightarrow \otimes \Gamma$ such that the operation $\mathcal{P}(\Gamma_1, \otimes \Gamma, \Gamma_2; \Delta) \rightarrow \mathcal{P}(\Gamma_1, \Gamma, \Gamma_2; \Delta)$ of precomposition with m_Γ is invertible. Dually, for any list of objects Δ , a *par product* (or *cotensor product*) of Δ is an object $\wp \Delta$ equipped with a polymap $w_\Delta: \wp \Delta \rightarrow \Delta$ such that the operation $\mathcal{P}(\Gamma; \Delta_1, \wp \Delta, \Delta_2) \rightarrow \mathcal{P}(\Gamma; \Delta_1, \Delta, \Delta_2)$ of postcomposition with w_Δ is invertible.

Definition 2.5 A *representable polycategory* is a polycategory that has tensors and pars of any finite lists of objects.

The definition of representable polycategory (called *two-tensor-polycategory* in [CS97]) may be alternatively stated requiring only the existence of binary and nullary tensors and pars, this being equivalent since the binary and nullary cases are sufficient for building up tensor and pars of arbitrary finite lists of objects. In any case, the definition implies that polymaps $\Gamma \rightarrow \Delta$ of a representable polycategory are in one-to-one correspondence with unary maps $\otimes \Gamma \rightarrow \mathfrak{Y} \Delta$ of its underlying category. Conversely, Cockett and Seely proved that any *linearly distributive category* $(\mathcal{C}, \otimes, 1, \mathfrak{Y}, \perp)$ induces a polycategory where the polymaps $\Gamma \rightarrow \Delta$ are defined as maps $\otimes \Gamma \rightarrow \mathfrak{Y} \Delta$ in \mathcal{C} , and that this extends to an equivalence of 2-categories between representable polycategories and linearly distributive categories [CS97]. One obtains $*$ -autonomous categories by moreover asking for the existence of *duals*.

Definition 2.6 A *right dual* of an object A is an object A^* equipped with polymaps $rcup_A : \cdot \rightarrow A, A^*$ and $rcap_A : A^*, A \rightarrow \cdot$ such that $rcup_A \circ_{A^*} rcap_A = id_A$ and $rcap_A \circ_A rcup_A = id_{A^*}$. A *left dual* of A is an object *A equipped with polymaps $lcup_A : \cdot \rightarrow {}^*A, A$ and $lcap_A : A, {}^*A \rightarrow \cdot$ such that $lcup_A \circ_{{}^*A} lcap_A = id_A$ and $lcap_A \circ_A lcup_A = id_{{}^*A}$.

Definition 2.7 A polycategory is said to *have duals* if any object has a right and a left dual.

Note that this definition may be simplified in the case of a symmetric polycategory because left and right duals coincide in that case, although following Cockett and Seely we have chosen to consider the more general situation. Cockett and Seely proved that in the symmetric case, representable polycategories with duals coincides with Barr’s notion of $*$ -autonomous categories [Bar91], and that in the non-symmetric case it coincides with a natural notion of “planar” $*$ -autonomous categories [CS97].


2.3 Representable polycategories with duals are $*$ -representable polycategories

In this section we introduce a notion of “ $*$ -representability” of a polycategory, and prove that a polycategory is $*$ -representable if and only if it is a representable polycategory with duals.

Definition 2.8 A polymap $u : \Gamma \rightarrow \Delta_1, A, \Delta_2$ is said to be *universal in the output A* (or *out-universal* for short, or simply *universal* when there is no ambiguity), written $u : \Gamma \rightarrow \Delta_1, \underline{A}, \Delta_2$ if for any polymap $h : \Gamma_1, \Gamma, \Gamma_2 \rightarrow \Delta_1, \Delta, \Delta_2$ such that $\Gamma_i = \emptyset$ or $\Delta_i = \emptyset$, there is a unique polymap $h/u : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$ such that $h = h/u \circ_A u$.

Dually, a polymap $n : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$ is *universal in the input A* (or *in-universal*), written $n : \Gamma_1, \underline{A}, \Gamma_2 \rightarrow \Delta$ if for any polymap $h : \Gamma_1, \Gamma, \Gamma_2 \rightarrow \Delta_1, \Delta, \Delta_2$ such that $\Gamma_i = \emptyset$ or $\Delta_i = \emptyset$ there is a unique polymap $n \setminus h : \Gamma \rightarrow \Delta_1, A, \Delta_2$ such that $h = n \circ_A n \setminus h$.

Graphically, the definitions are summarized in the following diagram:



Remark 2.9 By extension, we say that A is an *out-universal object* (resp. *in-universal object*) with respect to the surrounding context $\Gamma \rightarrow \Delta_1, _, \Delta_2$ (resp. $\Gamma_1, _, \Gamma_2 \rightarrow \Delta$) if there is an out-universal polymap $\Gamma \rightarrow \Delta_1, \underline{A}, \Delta_2$ (resp. in-universal polymap $\Gamma_1, \underline{A}, \Gamma_2 \rightarrow \Delta$). For a fixed surrounding context, in-universal and out-universal objects are unique up to unique isomorphism.

Definition 2.10 A polycategory is said to be *$*$ -representable* if it has all in-universal and out-universal objects, that is, if for any $\Gamma, \Delta_1, \Delta_2$ there is an object A equipped with an out-universal polymap $\Gamma \rightarrow \Delta_1, \underline{A}, \Delta_2$, and similarly, for any $\Gamma_1, \Gamma_2, \Delta$ there is an object A equipped with an in-universal polymap $\Gamma_1, \underline{A}, \Gamma_2 \rightarrow \Delta$.

It may be argued that Definition 2.8 is a natural generalisation of the notion of *strong universal* multimap in a multicategory [Her00], and Definition 2.10 the natural generalisation of representability from multicategories to polycategories (*pace* Defn. 2.5). In Section 3, we will see that these concepts are special cases of more general fibrational concepts. Like strong universal multimaps in a multicategory, both in-universal and out-universal polymaps are closed under composition in an appropriate sense.

Proposition 2.11 *In-universal polymaps compose, in the sense that if $f : \Gamma_1, \underline{A}, \Gamma_2 \rightarrow \Delta_1, B, \Delta_2$ (in the notation of Definition 2.8) and $g : \Gamma'_1, \underline{B}, \Gamma'_2 \rightarrow \Delta'$, then $g \circ_B f : \Gamma'_1, \Gamma_1, \underline{A}, \Gamma_2, \Gamma'_2 \rightarrow \Delta_1, \Delta', \Delta_2$. Similarly,*

out-universal maps compose in the sense that if $f : \Gamma \rightarrow \Delta_1, \underline{B}, \Delta_2$ and $g : \Gamma'_1, B, \Gamma'_2 \rightarrow \Delta'_1, \underline{C}, \Delta'_2$, then $g \circ_B f : \Gamma'_1, \Gamma, \Gamma'_2 \rightarrow \Delta_1, \Delta'_1, \underline{C}, \Delta'_2, \Delta_2$.

Proof. As we will see later, this is a special case of Proposition 3.4. \square

An immediate consequence of these definitions is that tensor products can be considered as out-universal objects, and par products as in-universal objects.

Proposition 2.12 *An object $\otimes \Gamma$ equipped with a polymap $m : \Gamma \rightarrow \otimes \Gamma$ is a tensor product of Γ iff m is out-universal (in its unique output). Dually, an object $\wp \Delta$ equipped with a polymap $w : \wp \Delta \rightarrow \Delta$ is a par product of Δ iff w is in-universal (in its unique input).*


Somewhat more surprisingly, duals can also be characterised as either in-universal or out-universal objects.

Proposition 2.13 *Let A and A^* be objects of a polycategory \mathcal{P} . The following are equivalent:*


- (i) *there is an out-universal map $rcup_A : \cdot \rightarrow A, \underline{A^*}$*
- (ii) *there is an in-universal map $rcap_A : \underline{A^*}, A \rightarrow \cdot$*
- (iii) *there is an out-universal map $rcup_A : \cdot \rightarrow \underline{A}, A^*$*
- (iv) *there is an in-universal map $rcap_A : A^*, \underline{A} \rightarrow \cdot$*
- (v) *A^* is the right dual of A*

Proof. We prove $(i) \leftrightarrow (v)$, the other cases are similar.

Suppose that there is an out-universal map $rcup_A : \cdot \rightarrow A, \underline{A^*}$. Then it verifies the following factorisation property



In particular by taking $h = id_A$ we get for $rcap_A := g : A^*, A \rightarrow \cdot$:



That satisfies the snake identity (where composition take place) in A^* by definition.

We get the second snake identity by following equalities where we go from the first line to the second by introducing a snake identity in A^* and we get the last equality by the interchange law for composition:

$$\begin{array}{c}
 \begin{array}{ccc}
 \begin{array}{|c|} \hline rcup_A \\ \hline \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \\ \square \\ \text{---} \end{array} \\ \hline \end{array} & = & \begin{array}{|c|} \hline rcup_A \\ \hline \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \end{array} \\ \hline \end{array} \\
 \\
 = & \begin{array}{|c|} \hline rcup_A \\ \hline \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \end{array} \\ \hline \end{array} & = & \begin{array}{|c|} \hline rcup_A \\ \hline \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \\ \hline \end{array}
 \end{array}
 \end{array}$$


So $rcup_A$ can be factored through itself and id_{A^*} or it can be factored through itself and the snake in A . But universality of $rcup_A$ assure unicity of this factorisation. So we get the second snake identity.

Conversely if we have $rcup_A$ and $rcap_A$ satisfying the snake identities we want the factorisation property that for any $h : \Gamma \rightarrow A, \Delta$ there is a unique $g : \Gamma, A^* \rightarrow \Delta$ such that $h = g \circ_{A^*} rcup_A$. We can take $g = rcap_A \circ_A h$. This satisfies the equation by using the snake identity and an interchange law. To get unicity it suffices to notice that for any g such that $h = g \circ_{A^*} rcup_A$ we can post-compose by $rcap_A$ to get $rcap_A \circ_A h = rcap_A \circ_A g \circ_{A^*} rcup_A = g \circ_{A^*} rcap_A \circ_A rcup_A = g$. \square


Remark 2.14 There is of course a similar result for left duals with $lcup_A$ and $lcap_A$.

Theorem 2.15 \mathcal{P} is a representable polycategory with duals iff it is $*$ -representable.

Proof. The right to left direction follows by propositions 2.12 and 2.13. For the left to right direction we want to construct in-universal and out-universal objects for any contexts just using \otimes, \wp and $*$. Given contexts $\Gamma, \Delta_1, \Delta_2$ consider the object $A := \Delta_1^* \otimes \Gamma \otimes \Delta_2^*$ where $\Delta_1^* := B_{1,n_1}^* \otimes \dots \otimes B_{1,1}^*$ for $\Delta_1 = B_{1,1}, \dots, B_{1,n_1}$ and similarly for Δ_2^* . This object comes with the following polypmap, which is a composition of out-universal polymaps along their out-universal objects. So by proposition 2.11, it is out-universal.



Similarly, given $\Gamma_1, \Gamma_2, \Delta$ the object $A := \Gamma_1^* \wp \Gamma_2^* \Delta$ is in-universal with in-universal polypmap:



\square

2.4 Examples

Example 2.16 Any linearly distributive category \mathcal{C} gives a polycategory $\mathcal{P}(\mathcal{C})$ called its underlying polycategory. It has the same objects as \mathcal{C} and a polypmap $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ in $\mathcal{P}(\mathcal{C})$ is a map $f : A_1 \otimes \dots \otimes A_m \rightarrow B_1 \wp \dots \wp B_n$ in \mathcal{C} .

Example 2.17 In particular any monoidal category gives rises to a polycategory with the same objects and with polymaps $f : A_1 \otimes \dots \otimes A_m \rightarrow B_1 \otimes \dots \otimes B_n$.


Example 2.18 Although this example is trivial, we will see that it plays an important role in Section 3. The terminal polycategory $\mathbb{1}$ has one object $*$ and a unique arrow $s_{m,n} : *^m \rightarrow *^n$ for every arity m and co-arity n .

Example 2.19 Any category induces a polycategory with only unary maps. Conversely any polycategory has an underlying category obtained by forgetting about the non-unary maps.

Example 2.20 From any multicategory \mathcal{M} we can define two polycategories \mathcal{M}^+ and \mathcal{M}^- that have the same objects as \mathcal{M} . The polymaps of \mathcal{M}^+ have always exactly one output and correspond to multimaps in \mathcal{M} while the polymaps in \mathcal{M}^- have always exactly one input and correspond to multimaps in \mathcal{M} reversed. Conversely from any polycategory we get two multicategories by restricting to polymaps with exactly one output and (reversed) polymaps with exactly one input.

Example 2.21 There are polycategories **Vect** and **FVect** of vector spaces (resp. finite dimensional vector spaces) and polylinear maps. Both of these can be seen as the underlying polycategories of monoidal categories of vector spaces and linear maps. **FVect** is a representable polycategory with duals while **Vect** is representable but does not have duals in general. In fact the vector spaces that admit a dual are precisely the finite dimensional ones.

Example 2.22 Free polycategories give examples of polycategories which are *not* representable. Let a “poly-signature” Σ consist of a collection of types, together with for any finite lists of types Γ and Δ , a set of operations $\Sigma(\Gamma; \Delta)$. The *free polycategory* generated by Σ , denoted $\mathcal{P}(\Sigma)$, has types as objects, and polymaps given by planar oriented trees with a boundary of free edges, whose nodes are labelled by operations and whose edges are labelled by types subject to the constraints specified by the signature. For example, here is a depiction of the composite polymap $f \circ_A (g \circ_B f) : A, B, B \rightarrow A, B$ in the free polycategory generated by the signature containing a pair of types A and B and a pair of operations $f : A, B \rightarrow B$ and $g : B \rightarrow A, A$ (in the diagram, the edges are implicitly oriented from left to right):



In general, composition is performed by grafting two trees along an edge, while the identity on a type A is given by the trivial tree with no nodes and one oriented edge labelled A . Observe this polycategory is not representable, for example there is no polymap $A, A \rightarrow A \otimes A$.

Example 2.23 A one-object multicategory is commonly referred to as an *operad*, while a one-object polycategory is also known as a *dioperad* [Gan03]. For any polycategory \mathcal{P} and any object $A \in \mathcal{P}$ there is a dioperad called the *endomorphism dioperad of A*, denoted $End_{\mathcal{P}}(A)$, defined as the full subpolycategory of \mathcal{P} containing only the object A . It has one object and its polymaps correspond to polymaps $A, \dots, A \rightarrow A, \dots, A$ in \mathcal{P} .

2.5 Example of Banach spaces

In this example we focus on Banach spaces. Although the use of polycategories is new most of the results are standard. For conciseness we omitted most of the definitions and proofs here, although they are available in an appendix of the paper. The standard theory of Banach spaces can be found in [Rya02]. We will only consider finite dimensional Banach spaces but this can be extended to the general case by replacing $*$ -autonomous structures by linearly distributive ones. This allows us to skip the subtleties about completeness.

We fix a field $\mathbb{K} = \mathbb{R}, \mathbb{C}$. **FVect** is the polycategory of finite dimensional \mathbb{K} -vector spaces and \mathbb{K} -polylinear maps, where a polylinear map $A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ corresponds to a linear map $A_1 \otimes \dots \otimes A_m \rightarrow B_1 \otimes \dots \otimes B_n$.

For a polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ and elements $a_i \in A_i$ and $\varphi_j \in B_j^*$ we will write the scalar $(\varphi_1, \dots, \varphi_n)f(a_1, \dots, a_m) := (\varphi_1 \otimes \dots \otimes \varphi_n)(f(a_1 \otimes \dots \otimes a_m))$. Continuous linear maps between Banach spaces correspond to bounded maps. This can be generalised to polylinear maps.

Definition 2.24 A polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ between normed vector spaces $(A_i, \| - \|_{A_i})$ and $(B_j, \| - \|_{B_j})$ is *bounded* if $\exists K, \forall a_i \in A_i, \forall \varphi_j \in B_j^*, |(\varphi_1, \dots, \varphi_n)f(a_1, \dots, a_m)| \leq K \prod_{i,j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}$.

Proposition 2.25 A unary polymap $f : A \rightarrow B$ is bounded if it is bounded as a linear map.

The smaller such K defines a norm on f and f is contractive when its norm is smaller than 1.

Definition 2.26 A polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ between normed vector spaces $(A_i, \| - \|_{A_i})$ and $(B_j, \| - \|_{B_j})$ is *contractive* if $\forall a_i \in A_i, \forall \varphi_j \in B_j^*, (\varphi_1, \dots, \varphi_n)f(a_1 \otimes \dots \otimes a_m) \leq \prod_{i,j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}$.

Definition 2.27 There are polycategories:

- **Ban** of Banach spaces and bounded polylinear maps
- **FBan** of finite dimensional Banach spaces and bounded polylinear maps
- **Ban₁** of Banach spaces and contractive polylinear maps
- **FBan₁** of finite dimensional Banach spaces and contractive polylinear maps

For objects in any of those polycategories to be isomorphic they need to be isomorphic as vector spaces. $(A, \| - \|)$ and $(A, \| - \|')$ are isomorphic in **Ban** and **FBan** if $\exists K, K', \forall a \in A, K\|a\| \leq \|a\|' \leq K'\|a\|$. Such norms are called equivalent. Two Banach spaces are isomorphic in **Ban₁** and **FBan₁** if their norms are equal.

In particular, this means that **FBan** is not an interesting polycategory since all the norms on a given finite dimensional vector space are equivalent.

Proposition 2.28 **FBan** is equivalent to **FVect**.

On the other hand, **FBan₁** is a $*$ -representable polycategory that does not come from a compact closed category. It is one of the examples of $*$ -autonomous categories described in Barr’s original paper [Bar79]. This is proved by using a characterisation of a $*$ -autonomous category as a symmetric monoidal closed category where the canonical maps $A \rightarrow A^{**}$ are isomorphisms. In particular, the induced norm for the par is never discussed. We did not find any reference in the literature linking it to the well-known injective norm in the theory of Banach spaces.

Definition 2.29 Let $(A, \| - \|_A)$ and $(B, \| - \|_B)$ be two Banach spaces. The *projective norm* $A \otimes B$ and the *injective norm* $A \wp B$ are the norms defined on the vector space $A \otimes B$ by the following formulas:

$$\|u\|_{A \otimes B} := \inf_{u = \sum_i a_i \otimes b_i} \|a_i\|_A \|b_i\|_B \quad \|u\|_{A \wp B} := \sup_{\|\varphi\|_{A^*}, \|\psi\|_{B^*} \leq 1} |(\varphi \otimes \psi)(u)|$$

These norms are known to be extremal among the set of well-behaved norms that one can put on the tensor.

Definition 2.30 For Banach spaces $(A, \| - \|_A)$ and $(B, \| - \|_B)$, a norm $\| - \|$ on $A \otimes B$ is a *crossnorm* if $\forall a, b \in A \times B, \|a \otimes b\| \leq \|a\|_A \|b\|_B$ and $\forall \varphi, \psi \in A^* \otimes B^*, \|\varphi \otimes \psi\|' \leq \|\varphi\|_{A^*} \|\psi\|_{B^*}$ with $\| - \|'$ the dual norm.

Remark 2.31 It is equivalent to ask for equalities in the definition. A proof can be found in [Rya02].

Proposition 2.32 A norm is a crossnorm iff it makes $A, B \rightarrow A \otimes B$ and $A \otimes B \rightarrow A, B$ contractive.

The injective and projective norms are crossnorms. The following property of the injective and projective crossnorm made us consider the injective crossnorm as a potential candidate for interpreting the par, and was one of our original motivations for studying the notion of bifibration of polycategories developed in Section 3.

Proposition 2.33 Let $\| - \|$ be a crossnorm then for any $u \in A \otimes B$ we have $\|u\|_{A \wp B} \leq \|u\| \leq \|u\|_{A \otimes B}$

Theorem 2.34 **FBan₁** is a $*$ -representable polycategory with tensor, par and duality defined above.

Remark 2.35 More than just a model of classical MLL, **FBan₁** is a model of classical MALL. The additive connectives are given by the vector space $A \oplus B$ with the norms $\|(a, b)\|_1 := \sum_i \|a\|_A + \|b\|_B$ and $\|(a, b)\|_\infty := \max(\|a\|_A, \|b\|_B)$. These norms are extremal among the p -norms.


3 Bifibrations of polycategories

In this section we introduce a notion of bifibration of polycategories, and prove that a polycategory is a representable polycategory with duals just in case it is bifibred over $\mathbb{1}$. We find it convenient to begin by adapting some terminological and notational conventions from the study of type refinement systems [MZ15, MZ16].

3.1 Definitions

Definition 3.1 A *poly-refinement system* is defined as a (strict) functor of polycategories $p : \mathcal{E} \rightarrow \mathcal{B}$. Explicitly, p sends objects $R \in \mathcal{E}$ to objects $p(R) \in \mathcal{B}$ and polymaps $\psi : R_1, \dots, R_m \rightarrow S_1, \dots, S_n$ in \mathcal{E} to polymaps $p(\psi) : p(R_1), \dots, p(R_m) \rightarrow p(S_1), \dots, p(S_n)$ in \mathcal{B} in such a way that identities and composition are preserved strictly. We write $R \sqsubset A$ (pronounced “ R refines A ”) to indicate that $p(R) = A$, and extend this to lists of objects in the obvious way, writing $\Pi \sqsubset \Gamma$ to indicate that $\Pi = R_1, \dots, R_n$ and $\Gamma = A_1, \dots, A_n$ for some $R_1 \sqsubset A_1, \dots, R_n \sqsubset A_n$. Finally, we write $\psi : \Pi \xrightarrow{f} \Sigma$ to indicate that ψ is a polymap $\Pi \rightarrow \Sigma$ in \mathcal{E} such that $p(\psi) = f$, with the implied constraint that $f : \Gamma \rightarrow \Delta$ where $\Pi \sqsubset \Gamma$ and $\Sigma \sqsubset \Delta$.


Remark 3.2 We will draw poly-refinement systems vertically. The top diagram will be in \mathcal{E} and the bottom one in \mathcal{B} with objects and polymaps directly above their image, e.g. preservation of composition is given by:



Definition 3.3 Fix $p : \mathcal{E} \rightarrow \mathcal{B}$ a poly-refinement system and $\psi : \Pi_1, R, \Pi_2 \xRightarrow[g]{\Sigma} \Sigma$ a polymap in \mathcal{E} with $R \sqsubset A$. ψ is *in-cartesian in R* (relative to p), written $\psi : \Pi_1, \underline{R}, \Pi_2 \xRightarrow[g]{\Sigma} \Sigma$, if for any polymap $\xi : \Pi_1, \Pi, \Pi_2 \xRightarrow[g \circ_A f]{\Sigma_1, \Sigma, \Sigma_2}$, satisfying the usual planarity condition that either $\Pi_i = \emptyset$ or $\Sigma_i = \emptyset$ for each $i = 1, 2$, there exists a unique polymap $\psi \setminus \xi : \Pi \xRightarrow[f]{\Sigma_1, R, \Sigma_2} \Sigma$ such that $\xi = \psi \circ_R (\psi \setminus \xi)$.

Dually, $\varphi : \Pi \xRightarrow[f]{\Sigma_1, S, \Sigma_2} \Sigma$, with $S \sqsubset B$ is *out-cartesian in S* , written $\varphi : \Pi \xRightarrow[f]{\Sigma_1, S, \Sigma_2} \Sigma$, if for any polymap $\xi : \Pi_1, \Pi, \Pi_2 \xRightarrow[g \circ_B f]{\Sigma_1, \Sigma, \Sigma_2}$, satisfying the planarity condition that either $\Pi_i = \emptyset$ or $\Sigma_i = \emptyset$, there is a unique polymap $\xi / \varphi : \Pi_1, S, \Pi_2 \xRightarrow[g]{\Sigma} \Sigma$ such that $\xi = \xi / \varphi \circ_S \varphi$.

Graphically, the definitions are summarised by the following diagram:



Proposition 3.4 *In-cartesian polymaps compose, in the sense that if $\varphi : \Pi_1, \underline{R}, \Pi_2 \xRightarrow[g]{\Sigma_1, S, \Sigma_2} \Sigma$ and $\psi : \Pi'_1, \underline{S}, \Pi'_2 \xRightarrow[f]{\Sigma'} \Sigma'$ then $\psi \circ_S \varphi : \Pi'_1, \Pi_1, \underline{R}, \Pi_2, \Pi'_2 \xRightarrow[g \circ_B f]{\Sigma_1, \Sigma', \Sigma_2} \Sigma$. Similarly, out-cartesian maps compose in the sense that if $\varphi : \Pi \xRightarrow[g]{\Sigma_1, \underline{S}, \Sigma_2} \Sigma$ and $\psi : \Pi'_1, S, \Pi'_2 \xRightarrow[f]{\Sigma'_1, \underline{T}, \Sigma'_2} \Sigma'$ then $\psi \circ_S \varphi : \Pi'_1, \Pi, \Pi'_2 \xRightarrow[g \circ_B f]{\Sigma_1, \Sigma'_1, \underline{T}, \Sigma'_2, \Sigma_2} \Sigma$.*

Definition 3.5 A poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$ is said to be a *pull-fibration* if for any $f : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$ in \mathcal{B} and any $\Pi_1 \sqsubset \Gamma_1$, $\Pi_2 \sqsubset \Gamma_2$, and $\Sigma \sqsubset \Delta$ there is an object $\mathbf{pull}[f](\Pi_1 \sqcup \Pi_2; \Sigma) \sqsubset A$ together with an in-cartesian polymap $\Pi_1, \mathbf{pull}[f](\Pi_1 \sqcup \Pi_2; \Sigma), \Pi_2 \xRightarrow[f]{\Sigma} \Sigma$. Dually, p is said to be a *push-fibration* if for any $f : \Gamma \rightarrow \Delta_1, B, \Delta_2$ in \mathcal{B} and any $\Pi \sqsubset \Gamma$, $\Sigma_1 \sqsubset \Delta_1$, and $\Sigma_2 \sqsubset \Delta_2$ there is an object $\mathbf{push}\langle f \rangle(\Pi; \Sigma_1 \sqcup \Sigma_2) \sqsubset B$ together with an out-cartesian polymap $\Pi \xRightarrow[f]{\Sigma_1, \mathbf{push}\langle f \rangle(\Pi; \Sigma_1 \sqcup \Sigma_2), \Sigma_2} \Sigma$. Finally, p is said to be a *bifibration* if it is both a pull-fibration and a push-fibration.

Remark 3.6 When pulling along a map $f : A \rightarrow \Delta$ with only one input, we will write $\mathbf{pull}[f](\Sigma)$ as shorthand for $\mathbf{pull}[f](\cdot; \Sigma)$. Similarly when pushing along a map $f : \Gamma \rightarrow A$, we will write $\mathbf{push}\langle f \rangle(\Gamma)$ for $\mathbf{push}\langle f \rangle(\Gamma; \cdot)$.

3.2 **-autonomous categories as bifibrations of polycategories*

Comparing diagram (†) with diagram (*), the following statements are self-evident.

Proposition 3.7 *Let \mathcal{P} be a polycategory. A polymap $u : \Gamma \rightarrow \Delta_1, A, \Delta_2$ (resp. $u : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$) is out-universal (resp. in-universal) in A iff it is out-cartesian (resp. in-cartesian) with respect to the unique functor $\mathcal{P} \rightarrow \mathbb{1}$ into the terminal polycategory.*

Proposition 3.8 *\mathcal{P} is a *-representable polycategory iff $\mathcal{P} \rightarrow \mathbb{1}$ is a bifibration of polycategories.*

We then derive the following as a corollary of Theorem 2.15 and Cockett and Seely’s connection between *-autonomous categories and representable polycategories with duals.

Theorem 3.9 *There is an equivalence between planar *-autonomous categories and bifibrations over the terminal polycategory $\mathbb{1}$.*

This correspondence may be extended in a straightforward way to the case of ordinary (symmetric) *-autonomous categories by considering symmetric bifibrations, that is, symmetric poly-refinement systems (= functors of symmetric polycategories that strictly preserve identities, composition, and the symmetry actions) which are bifibrations in the above sense. We also expect that this result may be stated more precisely as an equivalence of 2-categories, but we leave this to future work.

One application of Theorem 3.9 is that it provides a way of decomposing a *-autonomous structure on a category, using elementary facts about cartesian polymaps.

Proposition 3.10 *For $p : \mathcal{P} \rightarrow \mathcal{E}$ and $q : \mathcal{E} \rightarrow \mathcal{B}$ poly-refinement systems, and $\psi : \Pi_1, R, \Pi_2 \xrightarrow{g} \Sigma$ a polymap in \mathcal{P} , if ψ is p -in-cartesian in $R \sqsubset A$ and g is q -in-cartesian in $A \sqsubset X$ then ψ is $q \circ p$ -in-cartesian in $R \sqsubset X$.*

Remark 3.11 Similarly, a p -out-cartesian polymap over a q -out-cartesian polymap is $(q \circ p)$ -out-cartesian.

Proposition 3.12 *Let $p : \mathcal{E} \rightarrow \mathcal{B}$ be a poly-refinement system, and suppose that \mathcal{B} is *-representable. If \mathcal{E} has all in-cartesian liftings of in-universal polymaps and all out-cartesian liftings of out-universal polymaps then \mathcal{E} is a *-representable polycategory.*

Proof. By Propositions 3.7 and 3.10. □

3.3 *Additional examples*

Example 3.13 Let \mathcal{E} and \mathcal{B} be ordinary categories considered as degenerate polycategories with only unary co-unary maps (i.e., polymaps of arity and co-arity 1), and let $p : \mathcal{E} \rightarrow \mathcal{B}$ be an ordinary (strict) functor. Then p is a pull-fibration, push-fibration or bifibration just in case it is an ordinary (Grothendieck) fibration, opfibration or bifibration. Similarly, if \mathcal{E} and \mathcal{B} are multicategories considered as polycategories with only co-unary maps, then p is a push-fibration just in case it is a covariant fibration of multicategories in the sense of Hermida [Her04], and more generally the polycategorical notions of pullback and pushforward coincide with the multicategorical ones described in [Hĭ7,LSR17].

Example 3.14 The forgetful functor $\mathbf{Cat}_* \rightarrow \mathbf{Cat}$ from the category of pointed (small) categories to the category of (small) categories is an opfibration of 2-categories. The pushforward of (\mathcal{A}, A) along $F : \mathcal{A} \rightarrow \mathcal{B}$ is $(\mathcal{B}, F(A))$. Similarly the forgetful functor $\mathbf{Adj}_* \rightarrow \mathbf{Adj}$ of pointed adjunctions is a bifibration of 2-categories. Here a pointed adjunction between pointed categories (\mathcal{A}, A) and (\mathcal{B}, B) consist of an adjunction $F \dashv G : \mathcal{A} \rightarrow \mathcal{B}$ and a morphism $f : F(A) \rightarrow B$ in \mathcal{B} - or equivalently of a morphism $g : A \rightarrow G(B)$ in \mathcal{A} . The pushforward is given by the image by F while the pullback is given by the image of G . While working on the polycategorical Grothendieck correspondence we will define the (weak) 2-polycategory of multivariable adjunction \mathbf{MAdj} . It also has a pointed variant \mathbf{MAdj}_* . The forgetful functor induced is a bifibration of 2-polycategories.

3.4 *Forgetful functor from Banach spaces*

We will use proposition 3.10 to derive the *-representability of the polycategory \mathbf{FBan}_1 defined in 2.5. In order to do that we consider the forgetful functor $\mathbf{FBan}_1 \rightarrow \mathbf{FVect}$. We want to characterise the polymaps that admit cartesian liftings. The proofs can be found in appendix A.

Definition 3.15 Given $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ and norms $\| - \|_{A_i}, \| - \|_{B_j}$ for all $i \neq k$ and all j , we define a function $\| - \|_f : A_k \rightarrow \mathbb{K}$ by $\|x\|_f := \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m)|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}}$

Proposition 3.16 $\| - \|_f$ is a pseudonorm on A_k .

We want to characterise the polymaps for which this is a norm.

Definition 3.17 f is *injective in A_k* - or *A_k -injective* - if $(\forall a_i, f(a_1, \dots, x, \dots, a_m) = 0) \Rightarrow x = 0$

Definition 3.18 The A_k -kernel of f is the set $\text{Ker}_{A_k}(f) := \{x \in A_k \mid f(a_1, \dots, x, \dots, a_m) = 0 \forall a_i\}$.

The A_k -kernel of f forms a vector space. f is A_k -injective if its A_k -kernel is trivial. Furthermore we have that $\text{Ker}_{A_k}(f) = \{x \in A_k \mid (\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m) = 0 \forall a_i \forall \varphi_j\}$ by linearity of $\bigotimes_j \varphi_j$.

Remark 3.19 A polylinear map $f : A \rightarrow B$ is A -injective if it is injective as a linear map.

Proposition 3.20 For f , $\| - \|_{A_i}$ and $\| - \|_{B_j}$, $\| - \|_f$ is a norm iff f is A_k -injective.

It is worth noticing that this only depends on f and not on any properties of the norms.

Proposition 3.21 For f A_k -injective and norms $\| - \|_{A_i}, \| - \|_{B_j}$, the norm $\| - \|_f$ makes f contractive.

This norm defines a pullback in $\mathbf{Ban}_1, \mathbf{FBan}_1$.

Proposition 3.22 Given a B -injective polylinear map $g : \Gamma'_1, A, \Gamma'_2 \rightarrow \Delta'$ with lists $\Gamma'_i = A'_{i,1}, \dots, A'_{i,m'_i}$ and $\Delta' = B'_1, \dots, B'_n$ we fix families of norms $\| - \|_{\Gamma'_i} = (\| - \|_{A'_{i,j}})$ and $\| - \|_{\Delta'} = (\| - \|_{B'_i})$. Then the pullback is given by $\mathbf{pull}[g](\Gamma'_1, \| - \|_{\Gamma'_1} \smile (\Gamma'_2, \| - \|_{\Gamma'_2}); (\Delta, \| - \|_{\Delta})) = (A, \| - \|_g)$.

So we have in-cartesian liftings of any polylinear map that is injective in the input considered. The injectivity condition is only needed for $\| - \|_f$ to be a norm, otherwise it is still a seminorm, i.e., $\|x\|_f \geq 0$ for all x and $\|0\|_f = 0$, but $\|x\|_f = 0$ does not imply $x = 0$.

Corollary 3.23 There is a polycategory $\mathbf{FBan}_1^{\text{ps}}$ of finite dimensional complete seminormed vector spaces and contractive polylinear maps that comes with a forgetful functor that is pull-fibred.

Now we want to determine which polylinear maps have out-cartesian liftings.

Definition 3.24 For $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ and families of norms $\| - \|_{\Gamma}, \| - \|_{\Delta_1}, \| - \|_{\Delta_2}$, we define a function $\| - \|_f : B_k \rightarrow \bar{\mathbb{K}}$ where $\bar{\mathbb{K}}$ is the completion of \mathbb{K} , i.e., we add a point at infinity. It is given by $\|y\|_f := \inf_{y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)} \sum_i \|\vec{\varphi}_{1,i}\| \|\vec{\varphi}_{2,i}\| \|\vec{a}_i\|$ where the sum is over all the decompositions of y .

Proposition 3.25 $\| - \|_f$ is an extended norm, i.e., a norm with value in $\bar{\mathbb{K}}$.

Definition 3.26 $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ is A -surjective if $\forall y \in A, \exists \vec{\varphi}_{1,i}, \vec{\varphi}_{2,i}, \vec{a}_i, y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)$. The A -image of f is the set $\text{Im}_A(f) := \{\sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)\}$.

Proposition 3.27 $\text{Im}_A(f)$ forms a vector space. f is A -surjective iff $\text{Im}_A(f) = A$.

Remark 3.28 A linear map is B -surjective iff it is surjective. Indeed if for $y \in B$ there are x_i such that $y = \sum_i f(x_i)$ then by linearity $y = f(\sum_i x_i)$.

Proposition 3.29 For f and families of norms $\| - \|_{\Gamma}, \| - \|_{\Delta_1}, \| - \|_{\Delta_2}$, $\| - \|_f$ is a norm iff f is A -surjective.

Proposition 3.30 For f A -surjective and families of norms as usual, $\| - \|_f$ makes f contractive.

This norm defines a pushforward on $\mathbf{Ban}_1, \mathbf{FBan}_1$.

Proposition 3.31 For $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ a A -surjective polylinear map and the usual families of norms, we get the pushforward $\mathbf{push}\langle f \rangle(\Gamma; \Delta_1 \smile \Delta_2) = (A, \| - \|_f)$.

So we can take the out-cartesian lifting of any polymap that is surjective in the considered output.

Corollary 3.32 There are polycategories $\mathbf{FBan}_1^{\text{ex}}$ and $\mathbf{FBan}_1^{\text{ex,ps}}$ of f.d. extended normed/seminormed vector spaces and polylinear maps with forgetful functors that are push-fibred and bifibred respectively.

When considering \mathbf{FBan}_1 even without semi-/extended norms, there are still enough cartesian polymaps to lift the $*$ -representability of \mathbf{FVect} .

Proposition 3.33 *In \mathbf{FVect} , the universal polylinear maps $m_{A,B} : A, B \rightarrow A \otimes B$, $w_{A,B} : A \otimes B \rightarrow A, B$ and $rcap_A : A^*, A \rightarrow \cdot$ are $A \otimes B$ -surjective, $A \otimes B$ -injective and A^* -injective.*

Corollary 3.34 *\mathbf{FBan}_1 is $*$ -representable.*

Remark 3.35 We get the projective, injective and dual norm using the norms above: $\| - \|_{A \otimes B} = \| - \|_{m_{A,B}}$, $\| - \|_{A \otimes B} = \| - \|_{w_{A,B}}$ and $\| - \|_{A^*} = \| - \|_{rcap_A}$. The fact that the projective and injective crossnorms are extremal follows directly from the factorisation properties of the cartesian polymaps $m_{A,B}$ and $w_{A,B}$.

3.5 Frobenius monoids

Definition 3.36 In a polycategory \mathcal{P} a *Frobenius monoid* is an object A equipped with a unique polymap $\overline{(m, n)}_A : A^m \rightarrow A^n$ for each $m, n \in (N)$ such that $\overline{(1, 1)}_A = id_A$ and these polymaps are stable under composition.

Proposition 3.37 *Equivalently a Frobenius monoid in \mathcal{P} is a functor $F : \mathbb{1} \rightarrow \mathcal{P}$.*

Proof. The Frobenius monoid corresponds to $F(*)$ and the polymaps $\overline{(m, n)}_{F(*)}$ to $F(\overline{(m, n)})$. The properties needed on the polymaps are exactly functoriality of F . \square

Remark 3.38 For \mathcal{P} representable with $\otimes = \wp$ this reduces to the unbiased definition of a Frobenius monoid in a monoidal category.

Definition 3.39 Given a poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$ and a Frobenius monoid A in \mathcal{B} the *polyfiber of p over A* , noted $p^{-1}(A)$ is the subcategory of \mathcal{E} whose objects and polymaps are sent by p to A and the $\overline{(m, n)}_A$.

Proposition 3.40 *$p^{-1}(A)$ is equivalent to the following pullback:*

$$\begin{array}{ccc} p^{-1}(A) & \hookrightarrow & \mathcal{E} \\ \downarrow ! & \lrcorner & \downarrow p \\ \mathbb{1} & \xrightarrow{A} & \mathcal{B} \end{array} \quad \text{where } A : \mathbb{1} \rightarrow \mathcal{B} \text{ is the functor associated to the object } A.$$

Proposition 3.41 *Given a poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$ and a functor $s : \mathcal{B}' \rightarrow \mathcal{B}$, let $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$ be the pullback.*

$$\begin{array}{ccc} \mathcal{E} \times_{\mathcal{B}} \mathcal{B}' & \xrightarrow{\pi_1} & \mathcal{E} \\ \pi_2 \downarrow & \lrcorner & \downarrow p \\ \mathcal{B}' & \xrightarrow{s} & \mathcal{B} \end{array}$$

For a polymap $f : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$ in \mathcal{B}' and lists of objects Π_1, Π_2, Σ in $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$ lying over Γ_1, Γ_2 and Δ , if there is a pullback $pull_{s(f)}^{s(A)}(\pi_1(\Pi_1) | \pi_1(\Pi_2); \pi_1(\Sigma))$ in \mathcal{E} then there is a pullback $pull_f^A(\Pi_1 | \Pi_2; \Sigma)$ in $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$.

Proof. $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$ is the polycategory whose objects are pairs of objects (E, B') of \mathcal{E} and \mathcal{B} such that $p(E) = s(B')$ and whose polymaps are pairs of polymaps (f, b') such that $p(f) = s(b')$.

Given a polymap $f : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$ in \mathcal{B}' and lists of objects $(\Pi_1, \Gamma_1), (\Pi_2, \Gamma_2), (\Sigma, \Delta)$ in $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$ from a pullback $pull_{s(f)}^{s(A)}(\Pi_1 | \Pi_2; \Sigma)$ in \mathcal{E} with in-cartesian polymap $\varphi : \Pi_1, pull_{s(f)}^{s(A)}(\Pi_1 | \Pi_2; \Sigma), \Pi_2 \rightarrow \Sigma$ we get a pullback $pull_f^A((\Pi_1, \Gamma_1) | (\Pi_2, \Gamma_2); (\Sigma, \Delta)) := (pull_{s(f)}^{s(A)}(\Pi_1 | \Pi_2; \Sigma), A)$ with in-cartesian polymap (φ, f) . \square

Remark 3.42 Similarly if the pushforward exists in \mathcal{E} it exists in $\mathcal{E} \times_{\mathcal{B}} \mathcal{B}'$.

Corollary 3.43 *Given a poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$ and a Frobenius monoid $(A, \{\overline{(m, n)}_A\})$ in \mathcal{B} if all in-cartesian and out-cartesian polymaps of $\overline{(m, n)}_A$ exist then $p^{-1}(A)$ is $*$ -representable.*

4 Grothendieck correspondences

We should emphasise that the results in this section, and in particular the polycategorical Grothendieck correspondences, are conditioned on having a theory of weak 2-polycategories. To the extent of our knowledge such a theory has not been carefully worked out yet. We leave it as future work to craft this theory. Meanwhile we will describe the properties we assume to hold for weak 2-polycategories after recalling the usual notion of categorical Grothendieck correspondences.

4.1 Categorical Grothendieck correspondences

By ‘‘Grothendieck correspondence’’, we refer to the equivalence between fibrations and indexed categories, as well as a range of several other similar correspondences (the first was originally described by Bénabou [B 00]):

- Functor $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ lax normal functor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{Dist}$
- Fibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{Cat}$
- Opfibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{Cat}^{\text{op}}$ (or equivalently $\mathcal{B} \rightarrow \mathbf{Cat}$)
- Bifibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{Adj}$

Here \mathbf{Dist} is the bicategory whose objects are small categories and 1-cells $A \rightarrow B$ are *distributors*, i.e., functors $A \times B^{\text{op}} \rightarrow \mathbf{Set}$, with 2-cells given by natural transformations. While \mathbf{Cat} (respectively \mathbf{Adj}) is the (strict) 2-category whose objects are small categories, 1-cells are functors (resp. adjunctions), and 2-cells are natural transformations. Notice that \mathbf{Cat} , \mathbf{Cat}^{op} and \mathbf{Adj} are all subcategories of \mathbf{Dist} corresponding to distributors $A \rightarrow B$ that are representable in A , in B , and in both, respectively. The explicit constructions are provided in Appendix B.

4.2 Polycategorical Grothendieck correspondences

We want to extend the previous correspondences to polycategories as follows:

- Poly-refinement system $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ lax normal functor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{Dist}$
- Bifibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ Pseudofunctor $\mathcal{B}^{\text{op}} \rightarrow \mathbf{MAAdj}$

where \mathbf{Dist} is the weak 2-polycategory of sets and multivariable distributors and \mathbf{MAAdj} is the weak 2-polycategory of sets and multivariable adjunctions. It is worth noting that for us $(0, 0)$ -adjunctions will be sets, in contrast to the original definition in [Shu20] where they are taken to be trivial. Shulman discusses both possibilities, but chooses the latter to turn \mathbf{MAAdj} into a strict 2-polycategory, whereas the former fits more naturally in our framework at the price of \mathbf{MAAdj} being a weak 2-polycategory. Like in the categorical case, \mathbf{MAAdj} is a sub-2-polycategory of \mathbf{Dist} consisting of distributors that are representable in each of their variables.

4.2.1 About 2-polycategories

As stated above, to express these correspondences we need some theory of weak 2-polycategory, where by 2-polycategory we mean that the 1-cells can have multiple inputs and outputs but not the 2-cells. In this paper we only assume that there are weak 2-polycategories \mathbf{Dist} and \mathbf{MAAdj} and that lax functors and pseudofunctors behave in the expected way. More generally we suspect that weak 2-polycategories and $*$ -autonomous bicategories will be connected in a way such that the results of this paper can be relaxed to this setting. In particular, any compact closed bicategory – as defined by Mike Stay in [Sta16] – should be a $*$ -autonomous bicategory, and by extension a weak 2-polycategory. This would entail that \mathbf{Dist} is weak 2-polycategory.

4.2.2 Distributors and multivariable adjunctions

In this section we introduce the weak 2-polycategories \mathbf{Dist} and \mathbf{MAAdj} . We prove that a multivariable adjunction can be understood as a representable distributor.

Definition 4.1 \mathbf{Dist} is the weak 2-polycategory that has as objects categories, that has as polymaps $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ distributors $f : A_1 \times \dots \times A_m \rightarrow B_1 \times \dots \times B_n$ and that has as 2-cells natural transformations.

Definition 4.2 Given categories $A_1, \dots, A_m, B_1, \dots, B_n$, a (m, n) -adjunction or *multivariable adjunction* $(F_l)_{1 \leq l \leq n} \dashv (G_k)_{1 \leq k \leq m} : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ consists of the following data:

- functors $F_l : \prod_i A_i \times \prod_{j \neq l} B_j^{\text{op}} \rightarrow B_l$ for each l

- functors $G_k : \prod_{i \neq k} A_i^{\text{op}} \times \prod_j B_j \rightarrow A_k$ for each k
- natural isomorphisms $B_l(F_l(a_1, \dots, a_m, b_1, \dots, b_n), b_l) \simeq A_k(a_k, G_k(a_1, \dots, a_m, b_1, \dots, b_n))$ for any k, l

Example 4.3 A (1,1)-adjunction between A, B is a pair of functor $F : A \rightarrow B$ and $G : B \rightarrow A$ such that $B(F(a), b) = A(a, G(b))$. It is just a usual adjunction.

Example 4.4 Let $(\mathcal{C}, \otimes, I)$ be a biclosed monoidal category. By definition $(A \otimes -)$ has a right adjoint $A \multimap -$ and $(- \otimes B)$ has a right adjoint $- \multimap B$. We get three functors $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, $\multimap : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$ and $\circ : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$ such that $\mathcal{C}(A \otimes B, C) \simeq \mathcal{C}(B, A \multimap C) \simeq A, \mathcal{C} \circ B$, i.e. a (2,1)-adjunction $(\otimes) \dashv (\multimap, \circ)$.

Proposition 4.5 A (m, n) -adjunction $(F_l)_{1 \leq l \leq n} \dashv (G_k)_{1 \leq k \leq m} : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ is the same thing as a distributor $P : A_1 \times \dots \times A_m \rightarrow B_1 \times \dots \times B_n$ that is representable in each of its variables.


Proof. From any of the F_l we can define a distributor $P_l : A_1 \times \dots \times A_m \rightarrow B_1 \times \dots \times B_n$ representable in B_l by $P_l(-, -) := B_l(F_l(-), -)$. Similarly we can get distributors representable in A_k from the functors G_k by $P^k(-, -) := A_k(-, G_k(-))$. But all of these distributors are naturally isomorphic by definition of a multivariable adjunction.

Conversely given a distributor $P : A_1 \times \dots \times A_m \rightarrow B_1 \times \dots \times B_n$, representability in the A_k and B_l produce functors G_k with natural isomorphisms $P(-, -) \simeq A_k(-, G_k(-))$ and functors F_l with natural isomorphisms $P(-, -) \simeq B_l(F_l(-), -)$. \square

4.2.3 Fibres of a poly-refinement system and distributors between them

In the following we fix a poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$. We define a lax normal functor $\partial p : \mathcal{B}^{\text{op}} \rightarrow \mathbf{Dist}$ by considering the fibres of p like in the categorical case. We will use the convention that for any $\Gamma = A_1, \dots, A_n$, $p^{-1}(\Gamma) := p^{-1}(A_1) \times \dots \times p^{-1}(A_n)$. ∂p assigns to each object its fibre $\partial p(B) := p^{-1}(B) = \{S \in \mathcal{E} \mid p(S) = B\}$. To a polymap $f : \Delta \rightarrow \Gamma$ in \mathcal{B}^{op} , that we will equivalently consider as a polymap $f : \Gamma \rightarrow \Delta$ in \mathcal{B} , is assigned a distributor between the fibres $\partial p(f) : p^{-1}(\Delta) \times p^{-1}(\Gamma)^{\text{op}} \rightarrow \mathbf{Set}$. This distributor consists of the set of polymaps lying over f acted on by pre- and post-composition. More precisely, given lists of objects in the fibres $\Pi = (R_1, \dots, R_m) \sqsubset \Gamma = (A_1, \dots, A_m), \Sigma = (S_1, \dots, S_n) \sqsubset \Delta = (B_1, \dots, B_n)$ we define the action of the distributor $\partial p(f)$ on these objects by $\partial p(f)(\Sigma, \Pi) := \{\varphi : \Pi \rightarrow \Sigma \mid p(\varphi) = f\}$. And given lists of polymaps in the fibre $\vec{\psi} = (\psi_i : R'_i \xrightarrow{id_{A_i}} R_i)_{1 \leq i \leq m}$ and $\vec{\xi} = (\xi_j : S_j \xrightarrow{id_{B_j}} S'_j)_{1 \leq j \leq n}$ we get $\partial p(f)(\vec{\xi}, \vec{\psi}) := \vec{\xi} \circ \circ \vec{\psi}$.

This can be represented graphically.




It can be noted that the polymaps in the fibre have one-object domain and codomain. This is because all the polymaps in the fibre lies over the identity polymap in the base.

This is summarised in the following definition.

Definition 4.6 For a poly-refinement system $p : \mathcal{E} \rightarrow \mathcal{B}$ we define the lax normal functor $\partial p : \mathcal{B}^{\text{op}} \rightarrow \mathbf{Dist}$ by:

- For any $B \in \mathcal{B}$, $\partial p(B) := \{S \in \mathcal{E} \mid p(S) = B\}$
- For any $f : \Gamma \rightarrow \Delta$ in \mathcal{B} , $\partial p(f) : p^{-1}(\Delta) \rightarrow p^{-1}(\Gamma)$ defined by:
 - For any $\Pi \sqsubset \Gamma, \Sigma \sqsubset \Delta$, $\partial p(f)(\Sigma, \Pi) := \{\varphi : \Pi \rightarrow \Sigma \mid p(\varphi) = f\}$
 - For any $\vec{\psi} = (\psi_i : R'_i \xrightarrow{id_{A_i}} R_i)_{1 \leq i \leq m}$ and $\vec{\xi} = (\xi_j : S_j \xrightarrow{id_{B_j}} S'_j)_{1 \leq j \leq n}$, $\partial p(f)(\vec{\xi}, \vec{\psi}) = \vec{\xi} \circ \circ \vec{\psi}$

The proof that this defines a lax normal functor is similar to the categorical one with some extra bookkeeping because of the presence of contexts of inputs and outputs. For example, given polymaps $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ and $g : \Gamma'_1, A, \Gamma'_2 \rightarrow \Delta'$ we define a natural transformation $\partial p(g) \circ_{\partial p(A)} \partial p(f) \Rightarrow \partial p(g \circ_A f)$ by the following


 Fig. 1. Polymap $\mathbf{pull}[f](\Pi_1 \smile \Pi_2; \Sigma) \rightarrow \mathbf{pull}[f](\Pi'_1 \smile \Pi'_2; \Sigma')$

family of multimaps:

$$\begin{aligned}
 (\partial p(g) \circ_{\partial p(A)} \partial p(f))(\Sigma_1, \Sigma', \Sigma_2, \Pi'_1, \Pi, \Pi'_2) &= \int^R \partial p(g)(\Sigma', \Pi'_1, R, \Pi'_2) \times \partial p(f)(\Sigma_1, R, \Sigma_2, \Pi) \\
 &= \int^R \{\varphi' : \Pi'_1, R, \Pi'_2 \rightarrow \Sigma' \mid p(\varphi') = g\} \times \{\varphi : \Pi \rightarrow \Sigma_1, R, \Sigma_2 \mid p(\varphi) = f\} \\
 &\rightarrow \{\varphi' : \Pi'_1, R, \Pi'_2 \rightarrow \Sigma' \mid p(\varphi') = g\} \times \{\varphi : \Pi \rightarrow \Sigma_1, R, \Sigma_2 \mid p(\varphi) = f\} \\
 &\rightarrow \{\varphi'' : \Pi'_1, \Pi, \Pi'_2 \rightarrow \Sigma_1, \Sigma', \Sigma'_2 \mid p(\varphi'') = g \circ_A f\}
 \end{aligned}$$

Now suppose that p is a bifibration and consider a polymap $f : \Gamma_1, A, \Gamma_2 \rightarrow \Delta$. We define a functor $\mathbf{pull}[f]_A : p^{-1}(\Delta) \times p^{-1}(\Gamma_1)^{\text{op}} \times p^{-1}(\Gamma_2)^{\text{op}} \rightarrow p^{-1}(A)$ by sending any $\Sigma \sqsubset \Delta, \Pi_i \sqsubset \Gamma_i$ to $\mathbf{pull}[f](\Pi_1 \smile \Pi_2; \Sigma)$. From lists of polymaps $\vec{\psi}_1, \vec{\psi}_2$ and $\vec{\xi}$ we get a polymap $\mathbf{pull}[f](\Pi_1 \smile \Pi_2; \Sigma) \rightarrow \mathbf{pull}[f](\Pi'_1 \smile \Pi'_2; \Sigma')$ by using the factorisation property of $\mathbf{pull}[f](\Pi'_1 \smile \Pi'_2; \Sigma')$. It is represented in Figure 1 where the two big blank boxes are the in-cartesian polymaps associated to the pullbacks.

By the universal property of the pullback we can link $\partial p(f)$ and $\mathbf{pull}[f]_A$ in the following way.

$$\begin{aligned}
 \partial p(f)(\Sigma, \Pi_1, -, \Pi_2) &= \{\varphi : \Pi_1, -, \Pi_2 \rightarrow \Sigma \mid p(\varphi) = f\} \\
 &= \{\psi : - \rightarrow \mathbf{pull}[f](\Pi_1 \smile \Pi_2; \Sigma) \mid p(\psi) = id_A\} \\
 &= Hom_{p^{-1}(A)}(-, \mathbf{pull}[f]_A(\Sigma, \Pi_1, \Pi_2))
 \end{aligned}$$

This makes $\partial p(f)$ representable in A . Since by definition of a bifibration we get such a pull-functor for each of the inputs of f and some similar push-functors for the outputs this makes $\partial p(f)$ a multivariable adjunction. Since it is true for any polymap f in \mathcal{B}^{op} we get that ∂p factors through \mathbf{MAdj} . Finally the fact that cartesian polymaps compose makes $\partial p : \mathcal{B}^{\text{op}} \rightarrow \mathbf{MAdj}$ a pseudofunctor.

4.2.4 Polycategorical Grothendieck-Bénabou construction

Conversely, given a lax normal functor $F : \mathcal{B}^{\text{op}} \rightarrow \mathbf{Dist}$ we construct its polycategory of elements $\int F$.

Definition 4.7 The polycategory of elements $\int F$ has:

- for objects, pairs (A, R) with $A \in \mathcal{B}$ and $R \in F(A)$
- for polymaps $(f, \varphi) : (\Gamma, \Pi) \rightarrow (\Delta, \Sigma)$, pairs of a polymap $f : \Gamma \rightarrow \Delta$ in \mathcal{B} and an element $\varphi \in F(f)(\Sigma, \Pi)$
- for identities (id_A, id_R)

- for composition $(g, \psi) \circ_{(A,R)} (f, \varphi) = (g \circ_A f, \mu(\widetilde{(\varphi, \psi)}))$ where:
 - $\widetilde{(\varphi, \psi)} \in (F(g) \circ_{F(A)} F(f))(\Sigma_1, \Sigma', \Sigma_2, \Pi_1', \Pi, \Pi_2')$ is the canonical element induced by the elements $\varphi \in F(f)(\Sigma_1, R, \Sigma_2, \Pi)$ and $\psi \in F(g)(\Sigma', \Pi_1', R, \Pi_2')$
 - $\mu : F(g) \circ_{F(A)} F(f) \Rightarrow F(g \circ_A f)$ is the natural transformation giving lax functoriality of F

The fact that this is a polycategory follows from the coherence laws of F . Furthermore it can be proven that these constructions are inverse to each other using the same arguments that for the categorical constructions.

4.3 Frobenius pseudomonoids and Classical Linear Logic

Like in Section 3.5 there are different ways to define a Frobenius pseudomonoid. The most convenient in our case will be to think of those as (the image of) a pseudofunctor out of $\mathbb{1}$.

Definition 4.8 A *Frobenius pseudomonoid* in a 2-polycategory \mathcal{C} is a pseudofunctor $F : \mathbb{1} \rightarrow \mathcal{C}$.

Using the polycategorical Grothendieck correspondence we recover the result recently advertised by Shulman that Frobenius pseudomonoids in **MAAdj** are equivalent to $*$ -autonomous categories.

Theorem 4.9 (Shulman [Shu19]) *There is a correspondence between Frobenius pseudomonoid and $*$ -autonomous categories.*

Proof. Using the polycategorical Grothendieck correspondence, pseudofunctors $\mathbb{1} \rightarrow \mathbf{MAAdj}$ correspond to bifibrations $p : \mathcal{E} \rightarrow \mathbb{1}$. Then using theorem 3.9 these correspond to $*$ -representable polycategories. \square

Remark 4.10 Given a Frobenius monoid $(A, \overline{(m, n)}_A)$ in \mathcal{B} and a lax normal functor $F : \mathcal{B} \rightarrow \mathbf{Dist}$. The polyfibre of A relatively to the functor $\int F$ such as defined in Section 3.5 is given by the image of A (and the polymaps $\overline{(m, n)}_A$) by F . If F is pseudo on these polymaps this forms a Frobenius pseudomonoid in **Dist**. When the image of these polymaps are representable in all their variables this factors through **MAAdj** giving a $*$ -representable polycategory. This is the another way of understanding the result 3.43.

5 Conclusion and Further work

In this article we developed some of the theory of bifibred polycategories and provided examples of applications.

We started by recasting the notion of representable polycategory with duals as the equivalent notion of $*$ -representable polycategory, in a way that made it clear that it is a special kind of bifibred polycategory. We then explored examples of how this connection can be used to lift the connectives \otimes , \wp and $(-)^*$ from the codomain to the domain of a functor of polycategories if this functor has good fibrational properties. Since $*$ -representable polycategories are equivalent to $*$ -autonomous categories this helped us to analyse some models of classical MLL from a new perspective.

In the example of Banach spaces the base polycategory is induced by a compact closed category. Compact closed categories are known to provide good models for many different kinds of systems and processes, and have a simple graphical language of string diagrams. In future work we would like to explore in more detail examples of this sort. It would be interesting to study functors $p : \mathcal{E} \rightarrow \mathcal{B}$ where \mathcal{B} is the underlying polycategory of a compact closed category modelling systems and processes while \mathcal{E} is the underlying polycategory of a $*$ -autonomous category modelling properties of those systems and processes. An example that is closely related is the work on causal structures developed in [KU17]. In this paper the authors construct a $*$ -autonomous category of systems and causal processes out of any compact closed category of systems and processes with some specific discard maps. The $*$ -autonomous structure of the causal category can be lifted along the forgetful functor in the same way as we have done for Banach spaces. In fact these are instances of a generic construction that assigns to any compact closed category a $*$ -representable polycategory whose objects consist of objects of the original category together with a set of states (or global elements) with some closure condition. In the case of causal structures, the set corresponds to the causal states while in the case of the Banach spaces it corresponds to the (sub)normalised states, i.e., to the unit sphere (or ball). We can think of these sets as expressing some properties about the systems by specifying the states that satisfy it.

Multicategories are highly asymmetrical in their treatment of inputs and outputs. Polycategories restore this symmetry, which makes them an elegant object of study, but also a somewhat more complex object and more difficult to find in nature. Just as a $*$ -autonomous category can be decomposed into a pair of monoidal categories related by a monoidal duality, we believe it is worth trying to decompose the notion of bifibration of polycategories into a pair of fibrations of multicategories related by duality. Such an analysis would likely be in the spirit of Cockett and Seely's *polarized polycategories* [CS07] and Mellies' *chiralities* [Mel16], and perhaps related to the notion of *bifibration chirality* introduced in [MZ16].

Acknowledgements

We thank the anonymous MFPS reviewers, as well as Mike Shulman for several very helpful comments made during the online edition of MFPS 2020.

References

- [B00] Jean Bénabou. Distributors at work. Notes from a course at TU Darmstadt in June 2000, taken by Thomas Streicher, 2000.
- [Bar79] Michael Barr. **-Autonomous Categories*. Springer-Verlag, Berlin, Heidelberg, 1979.
- [Bar91] Michael Barr. *-autonomous categories and linear logic. *MSCS*, 1(2):159–178, 1991.
- [BCST96] R. F. Blute, J. R. B. Cockett, R. A. G. Seely, and T. H. Trimble. Natural deduction and coherence for weakly distributive categories. *JPAA*, 113(2):229–296, 1996.
- [BPS94] R. F. Blute, Prakash Panangaden, and R.A.G. Seely. Fock space: A model of linear exponential types, 1994. Revised version of MFPS 1993 paper, “Holomorphic models of exponential types in linear logic”.
- [CS97] J.R.B. Cockett and R.A.G. Seely. Weakly distributive categories. *JPAA*, 114(2):133–173, 1997.
- [CS07] J. R. B. Cockett and R. A. G. Seely. Polarized category theory, modules, and game semantics. *TAC*, 18(2):4–101, 2007.
- [Gan03] Wee Liang Gan. Koszul duality for dioperads. *Math. Res. Lett*, 10(1):109–124, 2003.
- [Gir87] Jean-Yves Girard. Linear logic. *TCS*, 50:1–102, 1987.
- [H17] Fritz Hörmann. Fibered multiderivators and (co)homological descent. *TAC*, 32(38):1258–1362, 2017.
- [Her00] Claudio Hermida. Representable multicategories. *Adv. Math.*, 151(2):164–225, 2000.
- [Her04] Claudio Hermida. Fibrations for abstract multicategories. In George Janelidze, Bodo Pareigis, and Walter Tholen, editors, *Galois Theory, Hopf Algebras, and Semiabelian Categories*, volume 43 of *Fields Institute Communications*. AMS, 2004.
- [Hyl02] J. M. E. Hyland. Proof theory in the abstract. *APAL*, 114(1-3):43–78, 2002.
- [KU17] Aleks Kissinger and Sander Uijlen. A categorical semantics for causal structure. In *Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS 2017)*, pages 1–12. IEEE Computer Society, June 2017.
- [Lam58] Joachim Lambek. The mathematics of sentence structure. *The American Mathematical Monthly*, 65(3):154–170, 1958.
- [Lam61] Joachim Lambek. On the calculus of syntactic types. In R. Jakobson, editor, *Structure of Language and its Mathematical Aspects*, Proceedings of Symposia in Applied Mathematics, XII, pages 166–178. AMS, 1961.
- [Lam69] Joachim Lambek. Deductive systems and categories II: Standard constructions and closed categories. In P. Hilton, editor, *Category Theory, Homology Theory and their Applications, I*, volume 86 of *LNM*, pages 76–122. Springer, 1969.
- [LSR17] Daniel R. Licata, Michael Shulman, and Mitchell Riley. A fibrational framework for substructural and modal logics. In *2nd International Conference on Formal Structures for Computation and Deduction (FSCD 2017)*, pages 25:1–25:22, 2017.
- [Mel16] Paul-André Melliès. Dialogue categories and chiralities. *Publications of the Research Institute for Mathematical Sciences*, 52, 01 2016.
- [MZ15] Paul-André Melliès and Noam Zeilberger. Functors are type refinement systems. In *Proceedings of the 42nd ACM SIGPLAN-SIGACT Symposium on Principles of Programming (POPL 2015)*, pages 3–16, January 2015.
- [MZ16] Paul-André Melliès and Noam Zeilberger. A bifibrational reconstruction of Lawvere’s presheaf hyperdoctrine. In *Proceedings of the 31st Annual IEEE Conference on Logic in Computer Science (LICS 2016)*, pages 555–564, July 2016.
- [Rya02] R.A. Ryan. *Introduction to Tensor Products of Banach Spaces*. Springer Monographs in Mathematics. Springer London, 2002.
- [Shu19] Michael Shulman. *-autonomous categories are Frobenius pseudomonoids. Referenced in [Shu20]. See also n-Category Café blog post (Nov. 17, 2017), “[Star-autonomous Categories are Pseudo Frobenius Algebras](#)”. In preparation, 2019.
- [Shu20] Michael Shulman. The 2-Chu-Dialectica construction and the polycategory of multivariable adjunctions. *TAC*, 35:89–136, 2020.
- [Sta16] Michael Stay. Compact Closed Bicategories. *TAC*, 31(26):755–798, 2016.
- [Sza75] M.E. Szabo. Polycategories. *Communications in Algebra*, 3(8):663–689, 1975.

A Application to Banach spaces

In this appendix we will develop enough basic theory of Banach spaces to make the statement above precise. Although the presentation using polycategories is new most of the results are standard and can be found in [Rya02].

In the following we will fix a field $\mathbb{K} = \mathbb{R}, \mathbb{C}$. **Vect** is the polycategory of \mathbb{K} -vector spaces and \mathbb{K} -polylinear maps, where a polylinear map $A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ correspond to a linear map $A_1 \otimes \dots \otimes A_m \rightarrow B_1 \otimes \dots \otimes B_n$. **FBVect** is its subpolycategory consisting of finite dimensional vector spaces.

For a polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ and elements $a_i \in A_i$ and $\varphi_j \in B_j^*$ we will write the scalar $(\varphi_1, \dots, \varphi_n)f(a_1, \dots, a_m) := (\varphi_1 \otimes \dots \otimes \varphi_n)(f(a_1 \otimes \dots \otimes a_m))$.

Definition A.1 For a vector space V a *pseudonorm* or *seminorm* on V is a function $\| - \| : V \rightarrow \mathbb{K}$ such that

- $\|0\| = 0$
- $\|\lambda v\| = |\lambda| \|v\|$ for $\lambda \in \mathbb{K}$ and $v \in V$
- $\|v + w\| \leq \|v\| + \|w\|$ for $v, w \in V$

A *norm* is a pseudonorm such that $\|v\| = 0 \Rightarrow v = 0$.

A vector space V is *complete* for a norm $\| - \|$ if for any sequences $(v_i) \in \mathbb{K}^{\mathbb{N}}$, $\sum_{i=1}^{\infty} \|v_i\| = 0$ implies that $\sum_{i=1}^{\infty} v_i$ converges.

A *Banach space* is a complete normed vector space.

Any norm on a vector space induces a distance $d(u, v) := \|u - v\|$ that defines a topology on V . So we can talk of continuous linear maps. For a linear map to be continuous it suffices that it is continuous in 0. Furthermore it is equivalent to ask for the linear map to be bounded, $\|f\| := \sup_{\|x\| \leq 1} \|f(x)\| < \infty$. We could

also ask for a scalar $K \in \mathbb{K}$ such that for any $x \in V$ we have $\|f(x)\| \leq K\|x\|$. More generally we will define a notion of bounded polylinear maps.

Definition A.2 A polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ between normed vector spaces $(A_i, \| - \|_{A_i})$ and $(B_j, \| - \|_{B_j})$ is *bounded* if there is K such that for any $a_i \in A_i$ and any $\varphi_j \in B_j^*$ we have $|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, a_m)| \leq K \prod_{i,j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}$.

Proposition A.3 A *unary polymap* $f : A \rightarrow B$ is bounded if it is bounded as a linear map.

Proof. Suppose that there is K such that for any $x \in A$ and $\varphi \in B^*$ we have $|\varphi(f(x))| \leq K\|x\|_A \|\varphi\|_{B^*}$. Using the Hahn-Banach theorem given $f(x) \in B$ we can find $\varphi \in B^*$ such that $\varphi(f(x)) = \|f(x)\|_B$ and $\|\varphi\|_{B^*} \leq 1$. So for this choice of φ we get $\|f(x)\|_B \leq K\|x\|_A$ which means that f is bounded.

Conversely suppose that there is K such that for any $x \in A$ $\|f(x)\|_B \leq K\|x\|_A$. Then for any $\varphi \in B^*$ we have $\|\varphi\|_{B^*} \|f(x)\|_B \leq K\|x\|_A \|\varphi\|_{B^*}$. But then by definition, $\|\varphi\|_{B^*} \geq \frac{|\varphi(f(x))|}{\|f(x)\|_B}$ for $x \neq 0$. Finally we get $|\varphi(f(x))| \leq K\|x\|_A \|\varphi\|_{B^*}$ for any $x \neq 0$. For $x = 0$ both sides are null. \square

Like in the linear case we can take the smaller such K to be the norm of f . Then a polylinear map is said to be *contractive* if its norms is smaller than 1.

Definition A.4 A polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ between normed vector spaces $(A_i, \| - \|_{A_i})$ and $(B_j, \| - \|_{B_j})$ is *contractive* if for any $a_i \in A_i$ and any $\varphi_j \in B_j^*$ the following inequation holds:

$$|(\varphi_1, \dots, \varphi_n)f(a_1 \otimes \dots \otimes a_m)| \leq \prod_{i,j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}$$

Bounded polylinear maps compose, so do contractive ones.

Definition A.5 We define the following polycategories:

- **Ban** has Banach spaces for objects and bounded polylinear map as polymaps
- **FBan** is the subpolycategory of **Ban** consisting of finite dimensional Banach spaces
- **Ban₁** has Banach spaces for objects and contractive polylinear map as polymaps
- **FBan₁** consists of finite dimensional Banach spaces and contractive polymaps

For objects in any of those polycategories to be isomorphic they need to be isomorphic as vector spaces. Two Banach spaces $(A, \|\cdot\|)$ and $(A, \|\cdot\|')$ are isomorphic in **Ban** and **FBan₁** if there are scalars K, K' such that for any $a \in A$ we have $K\|a\| \leq \|a\|' \leq K'\|a\|$. Such norms are called equivalent. They are equivalent in **Ban₁** and **FBan₁** if the norms are equal. In particular **FBan** is not an interesting polycategory.

Proposition A.6 **FBan** is equivalent to **FVect**.

Proof. Because all norms on a finite dimensional vector space are equivalent. \square

There is are forgetful functors from these polycategories to **Vect** or **FVect** respectively. With the exception of **FBan** \rightarrow **FVect** these are not bifibration. In the following we will focus on **Ban₁** and **FBan₁**. We want to understand what are cartesian liftings in these polycategories.

Definition A.7 Given a polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$, and norms $\|\cdot\|_{A_i}, \|\cdot\|_{B_j}$ on A_i, B_j for A_i expect A_k and all B_j , we define a function $\|\cdot\|_f^k : A_k \rightarrow \mathbb{K}$ as follow:

$$\|x\|_{A_k} := \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m)|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}}$$

Proposition A.8 $\|\cdot\|_f^k$ is a pseudonorm on A_k .

Proof. First we want to prove that $\|0\|_f^k = 0$. For that we will use linearity of the maps involved.

$$\begin{aligned} \|0\|_f^k &= \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, 0, \dots, a_m)|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}} \\ &= \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1 \otimes \dots \otimes \varphi_n)(f(a_1 \otimes \dots \otimes 0 \otimes \dots \otimes a_m))|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}} \\ &= \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1 \otimes \dots \otimes \varphi_n)(f(0))|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}} \\ &= \sup_{a_i, \varphi_j \neq 0} \frac{|(\varphi_1 \otimes \dots \otimes \varphi_n)(0)|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}} \\ &= \sup_{a_i, \varphi_j \neq 0} \frac{|0|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}} \\ &= \sup_{a_i, \varphi_j \neq 0} 0 \\ &= 0 \end{aligned}$$

Similarly using linearity of the maps and the fact that $|\cdot|$ is a norm we get that $\|\lambda x\|_f^k = |\lambda| \|x\|_f^k$ and $\|x + y\|_f^k \leq \|x\|_f^k + \|y\|_f^k$. \square

We want to characterise these polymaps for which this pseudonorm is a norm.

Definition A.9 A polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ is called *injective in A_k* or *A_k -injective* if the following condition is true $(\forall a_i, f(a_1, \dots, x, \dots, a_m) = 0) \Rightarrow x = 0$

Definition A.10 For a polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ its A_k -kernel is the set

$$Ker_{A_k}(f) := \{x \in A_k \mid f(a_1, \dots, x, \dots, a_m) = 0 \forall a_i\}$$

The A_k -kernel of a polylinear map forms a vector space. The polylinear map is A_k -injective if its A_k -kernel is trivial. Furthermore we have that $Ker_{A_k}(f) = \{x \in A_k \mid (\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m) = 0 \forall a_i\}$ by linearity of $\bigotimes_i \varphi_i$.

Remark A.11 For a unary map $f : A \rightarrow B$ to be A -injective means that it is injective as a linear map.

Proposition A.12 For a polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ and norms $\|-\|_{A_i}, \|-\|_{B_j}$ the pseudonorm $\|-\|_f^k$ is a norm iff f is A_k -injective.

Furthermore if A_k is infinite dimensional we note \widehat{A}_k its completion with respect to the norm $\|-\|_f^k$.

Proposition A.13 Given a A_k -injective polylinear map $f : A_1, \dots, A_m \rightarrow B_1, \dots, B_n$ and norms $\|-\|_{A_i}, \|-\|_{B_j}$, the norm $\|-\|_f^k$ makes f contractive.

Proof. By definition we have $\|a_k\|_f^k \geq \frac{|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m)|}{\prod_{i \neq k, j} \|a_i\|_{A_i} \|\varphi_j\|_{B_j^*}}$ for any a_i, φ_j . So $|(\varphi_1, \dots, \varphi_n)f(a_1, \dots, x, \dots, a_m)| \leq \prod_{i \neq k, j} \|a_i\|_{A_i} \|a_k\|_f^k \|\varphi\|_{A_j}$. But this is exactly what it means for f to be contractive. \square

This norm defines a pullback in $\mathbf{Ban}_1, \mathbf{FBan}_1$.

Proposition A.14 Given a B -injective polylinear map $g : \Gamma'_1, A, \Gamma'_2 \rightarrow \Delta'$ with $\Gamma'_i = A'_{i,1}, \dots, A'_{i,m'_i}$ and $\Delta' = B'_1, \dots, B'_n$ we fix families of norm $\|-\|_{\Gamma'_i} = (\|-\|_{A'_{i,j}})$ and $\|-\|_{\Delta}$ makes there respective vector spaces complete. Then $\mathbf{pull}[g](\Gamma'_1, \|-\|_{\Gamma'_1} \text{ } \Gamma'_2, \|-\|_{\Gamma'_2}); (\Delta, \|-\|_{\Delta}) = (\widehat{A}, \|-\|_g)$.

Proof.

Take a polylinear map $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ and families of norms $\|-\|_{\Gamma}, \|-\|_{\Delta_1}, \|-\|_{\Delta_2}$ that makes the corresponding vector spaces complete. Furthermore suppose that $g \circ_A f$ is contractive. We want to prove that f is contractive when A is equipped with the norm $\|-\|_g$.

Before starting the proof we should fix some notations. We will note $\vec{\varphi}_1$ a tuple of elements of Δ_1^* . Similarly we will have $\vec{\varphi}', \vec{\varphi}_2$. Finally we will have $\vec{a}'_1, \vec{a}, \vec{a}'_2$ for tuples of the Γ s. We will note $\|\vec{a}\|_{\Gamma} := \prod \|a_i\|_{A_i}$ and similarly for other tuples. Often we will omit the subscripts on the norms when they can be inferred from the context. Also notice that

$$(\vec{\varphi}_1, \vec{\varphi}', \vec{\varphi}_2)(g \circ_A f)(\vec{a}'_1, \vec{a}, \vec{a}'_2) = (\vec{\varphi}')g(\vec{a}'_1, (\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a}), \vec{a}'_2) \quad (\text{A.1})$$

where $(\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a}) := (\vec{\varphi}_1 \otimes id_A \otimes \vec{\varphi}_2) \circ f$ is a multilinear function into A .

To prove that f is contractive we need to prove that for any $\vec{\varphi}_1, \vec{\varphi}_2, \vec{a}$ and any $\alpha \in A^*$

$$|(\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})| \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\| \|\alpha\|'_g$$

where $\|-\|'_g$ is the norm dual to $\|-\|_g$. Without loss of generality we can take the tuples to not contain 0. By expanding the definition we need to prove

$$\frac{|(\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})|}{\|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\|} \leq \sup_{x \neq 0} \frac{|\alpha(x)|}{\|x\|_g}$$

We only need to find one x for which this inequality holds. Take $x = (\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a})$. We have that $\alpha(x) = (\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})$. So we just need to prove $\|(\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a})\|_g \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\|$. By expanding the definition of the norm we want $\sup \frac{|(\vec{\varphi}')g(\vec{a}'_1, (\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a}), \vec{a}'_2)|}{\|\vec{\varphi}'\| \|\vec{a}'_1\| \|\vec{a}'_2\|} \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\|$. Which means that we need to prove that for any $\vec{\varphi}_1, \vec{\varphi}', \vec{\varphi}_2, \vec{a}'_1, \vec{a}, \vec{a}'_2$ we have

$$|(\vec{\varphi}')g(\vec{a}'_1, (\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a}), \vec{a}'_2)| \leq \|\vec{\varphi}_1\| \|\vec{\varphi}'\| \|\vec{\varphi}_2\| \|\vec{a}'_1\| \|\vec{a}\| \|\vec{a}'_2\|$$

But then using A.1,

$$|(\vec{\varphi}_1, \vec{\varphi}', \vec{\varphi}_2)(g \circ_A f)(\vec{a}'_1, \vec{a}, \vec{a}'_2)| \leq \|\vec{\varphi}_1\| \|\vec{\varphi}'\| \|\vec{\varphi}_2\| \|\vec{a}'_1\| \|\vec{a}\| \|\vec{a}'_2\|$$

This is true because $g \circ_A f$ is contractive. \square

This means that we can take the in-cartesian lifting of any polymap that is injective in the input considered. This injectivity is only needed for $\|-\|_f$ to be a norm.

Corollary A.15 *There are polycategories $\mathbf{Ban}_1^{\text{ps}}$, $\mathbf{FBan}_1^{\text{ps}}$ of complete seminormed vector spaces and contractive polylinear maps that come with forgetful functors that are pull-fibred.*

Now we want to determine which polylinear maps have out-cartesian liftings.

Definition A.16 Given a polylinear map $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ and families of norms $\| - \|_{\Gamma}, \| - \|_{\Delta_1}, \| - \|_{\Delta_2}$, we define a function $\| - \|_f : B_k \rightarrow \overline{\mathbb{K}}$ where $\overline{\mathbb{K}}$ is the completion of \mathbb{K} , i.e. we add a point at the infinity.

$$\|y\|_f := \inf_{y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)} \sum_i \|\vec{\varphi}_{1,i}\| \|\vec{\varphi}_{2,i}\| \|\vec{a}_i\|$$

where the sum is done on all the possible decompositions of $y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)$.

Proposition A.17 $\| - \|_f$ is an extended norm, i.e. a norm with value in $\overline{\mathbb{K}}$.

Definition A.18 A polylinear map $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ is called *A-surjective* if for any $y \in A$ there exist $\vec{\varphi}_{1,i}, \vec{\varphi}_{2,i}, \vec{a}_i$ such that $y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)$.

The *A*-image of f is the set $Im_A(f) := \{\sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)\}$.

Proposition A.19 $Im_A(f)$ forms a vector space. f is *A-surjective* iff $Im_A(f) = A$.

Remark A.20 A linear map is *B-surjective* iff it is surjective. Indeed if for $y \in B$ there are x_i such that $y = \sum_i f(x_i)$ then by linearity $y = f(\sum_i x_i)$.

Proposition A.21 Given a polylinear map $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ and families of norms $\| - \|_{\Gamma}, \| - \|_{\Delta_1}, \| - \|_{\Delta_2}$, $\| - \|_f$ is a norm iff f is *A-surjective*.

Proof. The value of $\|y\|_f$ is non-finite iff we are taking the infimum over the empty set which means that y is not in the *A*-image of f . \square

Proposition A.22 For $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ a *A-surjective* polylinear map and the families of norms as usual, $\| - \|_f$ makes f contractive.

Proof. We want to prove that for any $\vec{\varphi}_1, \vec{\varphi}_2, \vec{a}$ and any $\alpha \in A^*$ $|(\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})| \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\| \|\alpha\|'$ where $\| - \|'$ is the norm dual to $\| - \|_f$. So we want $|(\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})| \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\| \sup_y \frac{|\alpha(y)|}{\|y\|_f}$. We just need to find a value of y that satisfy the following inequality: $|(\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})| \|y\|_f \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\| \|\alpha(y)\|$. Take $y = (\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a})$, by definition $\|(\vec{\varphi}_1, id_A, \vec{\varphi}_2)f(\vec{a})\|_f \leq \|\vec{\varphi}_1\| \|\vec{\varphi}_2\| \|\vec{a}\|$. Furthermore $\alpha(y) = (\vec{\varphi}_1, \alpha, \vec{\varphi}_2)f(\vec{a})$. So by multiplying on both side by $\alpha(y)$ we get the inequality that we wanted. \square

This norms define a pushforward on $\mathbf{Ban}_1, \mathbf{FBan}_1$.

Proposition A.23 For $f : \Gamma \rightarrow \Delta_1, A, \Delta_2$ a *A-surjective* polylinear map and the families of norms as usual, $\text{push}\langle f \rangle(\Gamma; \Delta_1 \dashv \Delta_2) = (\widehat{A}, \| - \|_f)$.

Proof. Suppose that $g : \Gamma'_1, A, \Gamma'_2 \rightarrow \Delta'$ is such that $g \circ_A f$ is contractive. Consider any $\vec{\varphi}', \vec{a}'_1, \vec{a}'_2$ and $y \in A$. Since f is *A-surjective* we can find a decomposition of $y = \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i)$ that reaches the infimum,

i.e. such that $\|y\|^f = \sum \|\vec{\varphi}_{1,i}\| \|\vec{\varphi}_{2,i}\| \|\vec{a}_i\|$. Then we have

$$\begin{aligned}
 |(\vec{\varphi}')g(\vec{a}'_1, y, \vec{a}'_2)| &= |(\vec{\varphi}')g(\vec{a}'_1, \sum_i (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i), \vec{a}'_2)| \\
 &= |\sum_i (\vec{\varphi}')g(\vec{a}'_1, (\vec{\varphi}_{1,i}, id_A, \vec{\varphi}_{2,i})f(\vec{a}_i), \vec{a}'_2)| \\
 &= |\sum_i (\vec{\varphi}_{1,i}, \vec{\varphi}', \vec{\varphi}_{2,i})(g \circ_A f)(\vec{a}'_1, \vec{a}_i, \vec{a}'_2)| \\
 &\leq \sum_i |(\vec{\varphi}_{1,i}, \vec{\varphi}', \vec{\varphi}_{2,i})(g \circ_A f)(\vec{a}'_1, \vec{a}_i, \vec{a}'_2)| \\
 &\leq \sum_i \|\vec{\varphi}_{1,i}\| \|\vec{\varphi}'\| \|\vec{\varphi}_{2,i}\| \|\vec{a}'_1\| \|\vec{a}_i\| \|\vec{a}'_2\| \\
 &= \|\vec{\varphi}'\| \|\vec{a}'_1\| \|\vec{a}'_2\| \sum_i \|\vec{\varphi}_{1,i}\| \|\vec{\varphi}_{2,i}\| \|\vec{a}_i\| \\
 &= \|\vec{\varphi}'\| \|\vec{a}'_1\| \|\vec{a}'_2\| \|y\|^f
 \end{aligned}$$

where we used in order, (1) definition of y , (2) linearity, (3) definition of the composition, (4) triangle inequality for the norm $|\cdot|$, (5) contractivity of $g \circ_A f$, (7) that the decomposition consider reach the infimum. This inequality is what is needed to prove that g is contractive. \square

This means that we can take the out-cartesian lifting of any polymap that is surjective in the considered output.

Corollary A.24 *There are polycategories $\mathbf{Ban}_1^{\text{ex}}$, $\mathbf{FBan}_1^{\text{ex}}$, $\mathbf{Ban}_1^{\text{ex,ps}}$, $\mathbf{FBan}_1^{\text{ex,ps}}$ of extended normed vector spaces and extended seminormed vector spaces that come with forgetful functors that are push-fibred and bifibred.*

Even when considering \mathbf{Ban}_1 , \mathbf{FBan}_1 there are enough cartesian polymaps to lift the representability and $*$ -representability of \mathbf{Vect} and \mathbf{FVect} to them.

Proposition A.25 *In \mathbf{Vect} , \mathbf{FVect} , the universal polylinear maps $A, B \rightarrow A \otimes B$ and $A \otimes B \rightarrow A, B$ are $A \otimes B$ -surjective and $A \otimes B$ -injective. In \mathbf{FVect} , the universal polylinear maps $A^*, A \rightarrow$ is A^* -injective.*

Proof. By definition of the tensor product any $u \in A \otimes B$ is a linear combination of elements from A and B , $u = \sum_i a_i \otimes b_i$. $A \otimes B$ -injectivity is trivial. Finally given $\varphi \in A^*$ if for all $a \in A$ $\varphi(a) = 0$ then $\varphi = 0$. \square

Corollary A.26 *The category of finite dimensional Banach spaces and contractive linear maps is $*$ -autonomous.*

The norm that we get on the dual is the usual dual norms. For the tensor and par we get back well-known norms.

Definition A.27 Let $(A, \|\cdot\|_A)$ and $(B, \|\cdot\|_B)$ be two Banach spaces.

The *projective crossnorm* is the norm defined as follow $\|u\|_{A \otimes B} := \inf_{u = \sum_i a_i \otimes b_i} \|a_i\|_A \|b_i\|_B$.

The *injective crossnorm* is the norm defined by $\|u\|_{A \otimes B} := \sup_{\|\varphi\|_{A^*}, \|\psi\|_{B^*} \leq 1} |(\varphi \otimes \psi)(u)|$

These are the norms that we get by taking the pushforward along $A, B \rightarrow A \otimes B$ and the pullback along $A \otimes B \rightarrow A, B$. So we deduce immediately that they are the norms that make (the completion of) $A \otimes B$ a tensor/par in \mathbf{FBan}_1 . These norms are known to be extremal into the set of well-behaved norms that we can put on the tensor product.

Definition A.28 Let $(A, \|\cdot\|_A)$ and $(B, \|\cdot\|_B)$ be two Banach spaces. A norm $\|\cdot\|$ on $A \otimes B$ is said to be a *crossnorm* if for any $a, b \in A \times B$ we have $\|a \otimes b\| \leq \|a\|_A \|b\|_B$ and for any $\varphi, \psi \in A^* \otimes B^*$ we have $\|\varphi \otimes \psi\|' \leq \|\varphi\|_{A^*} \|\psi\|_{B^*}$ with $\|\cdot\|'$ the norm dual to $\|\cdot\|$.

Remark A.29 It would have been equivalent to ask for equalities in the definition. A proof can be found in [Rya02].

Proposition A.30 *A norm on $A \otimes B$ is a crossnorm iff it makes the polylinear maps $A, B \rightarrow A \otimes B$ and $A \otimes B \rightarrow A, B$ contractive.*

Proof. By definition. \square

The injective and projective crossnorms are indeed crossnorms. The following property of the injective and projective crossnorm is known. It is this property that first make us consider the injective crossnorm as a potential candidate for interpreting the par. Getting a conceptual comprehension of why this intuition was true is what motivated the theory developed in this paper.

Proposition A.31 *Let $\| - \|$ be a crossnorm then for any $u \in A \otimes B$ we have $\|u\|_{A \otimes B} \leq \|u\| \leq \|u\|_{A \otimes B}$*

Proof. We have that $A, B \rightarrow A \otimes B$ factors through itself followed by $id_{A \otimes B}$. Since $\| - \|$ is a crossnorm $(A, \| - \|_A), (B, \| - \|_B) \rightarrow (A \otimes B, \| - \|)$ is contractive. But then since $(A \otimes B, \| - \|_{A \otimes B})$ is a pullback we can use its universal property to factor $(A, \| - \|_A), (B, \| - \|_B) \rightarrow (A \otimes B, \| - \|)$ through $(A, \| - \|_A), (B, \| - \|_B) \rightarrow (A \otimes B, \| - \|_{A \otimes B})$. This means that the identity map lift to a contractive map $(A \otimes B, \| - \|_{A \otimes B}) \rightarrow (A \otimes B, \| - \|)$. In other words for any $u \in A \otimes B$, $\|u\| \leq \|u\|_{A \otimes B}$. Similarly we can use the fact that $A \otimes B$ is a pushforward to get the other inequality. \square

Remark A.32 More than just a model of classical MLL (without negation), **FBan₁** (**Ban₁**) is a model of classical MALL (without negation). The additive connectives are given by the vector space $A \oplus B$ with the norms $\|(a, b)\|_1 := \sum_i \|a\|_A + \|b\|_B$ and $\|(a, b)\|_\infty := \max(\|a\|_A, \|b\|_B)$. These norms are extremal amongst the p -norms.

B Categorical Grothendieck correspondences

We recall a range of categorical Grothendieck correspondences that goes as follow:

- Functor $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ Lax normal functor $\mathcal{B} \rightarrow \mathbf{Dist}$
- Fibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B} \rightarrow \mathbf{Cat}$
- Opfibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B} \rightarrow \mathbf{Cat}^{\text{op}}$
- Bifibration $\mathcal{E} \rightarrow \mathcal{B} \longleftrightarrow$ pseudofunctor $\mathcal{B} \rightarrow \mathbf{Adj}$

Where **Dist** is the bicategory of (small) categories and distributors/profunctors and **Adj** is the 2-category of (small) categories and adjunctions. Notice that **Cat**, **Cat**^{op} and **Adj** are all subcategories of **Dist** corresponding to distributors $A \rightarrow B$ that are representable in A , in B , and in both, respectively.

Let us look at the details of this correspondence. We will first look at the left-to-right direction. Suppose that you have a functor $p : \mathcal{E} \rightarrow \mathcal{B}$ we want to get a lax normal functor $\partial p : \mathcal{B} \rightarrow \mathbf{Dist}$. We do that by considering the fibers over \mathcal{B} .

$$\begin{aligned}
 \partial p : \mathcal{B} &\rightarrow \mathbf{Dist} \\
 B &\mapsto p^{-1}(B) := \{E \in \mathcal{E} \mid p(E) = B\} \\
 \partial p(f) : p^{-1}(B)^{\text{op}} \times p^{-1}(B') &\rightarrow \mathbf{Set} \\
 f : B \rightarrow B' &\mapsto (E, E') \mapsto \{\varphi : E \rightarrow E' \mid p(\varphi) = f\} \\
 &\quad (\psi, \xi) \mapsto \xi \circ - \circ \psi
 \end{aligned}$$

We can see that this defines a lax normal functor.

$$\begin{aligned}
 (\partial p)(id_B)(E, E') &= \{\varphi : E \rightarrow E' \mid p(\varphi) = id_B\} = Hom_{p^{-1}(B)}(E, E') \\
 (\partial p)(id_B)(\psi, \xi) &= \xi \circ - \circ \varphi = Hom_{p^{-1}(B)}(\varphi, \xi)
 \end{aligned}$$

So $\partial p(id_B) = Hom_{p^{-1}(B)}$ which is the identity profunctor. We also want to find a natural transformation $(\partial p)(f') \circ (\partial p)(f) \Rightarrow (\partial p)(f' \circ f)$. By definition we have

$$\begin{aligned}
 ((\partial p)(f') \circ (\partial p)(f))(E, E'') &= \int^{E'} ((\partial p)(f') \times (\partial p)(f))(E, E'') \\
 &= \int^{E'} \{\varphi' : E' \rightarrow E'' \mid p(\varphi') = f'\} \times \{\varphi : E \rightarrow E' \mid p(\varphi) = f\}
 \end{aligned}$$

Then for any E' we get a function

$$\int^{E'} \{\varphi' : E' \rightarrow E'' \mid p(\varphi') = f'\} \times \{\varphi : E \rightarrow E' \mid p(\varphi) = f\} \rightarrow \{\varphi' : E' \rightarrow E'' \mid p(\varphi') = f'\} \times \{\varphi : E \rightarrow E' \mid p(\varphi) = f\}$$

And finally a function

$$\{\varphi' : E' \rightarrow E'' \mid p(\varphi') = f'\} \times \{\varphi : E \rightarrow E' \mid p(\varphi) = f\} \rightarrow \{\varphi'' : E \rightarrow E'' \mid p(\varphi'') = f' \circ f\}$$

by taking $\varphi'' = \varphi' \circ \varphi$ and using functoriality of p , $p(\varphi' \circ \varphi) = p(\varphi') \circ p(\varphi) = f' \circ f$.

The morphism that we get does not depend on the choice of E' by the universal property of the coend. We still need to prove the naturality condition and the coherence laws for a lax functor. This is left to the reader.

Now suppose that we want for ∂p to factor through $\mathbf{Cat} \hookrightarrow \mathbf{Dist}$. This means that for any $f : B \rightarrow B'$ we want a functor $p_*(f) : p^{-1}(B') \rightarrow p^{-1}(B)$ such that $\partial p(f) = \text{Hom}_{p^{-1}(B)}(-, p_*(-))$. If we unravel the definition

$$E \xrightarrow[f]{\varphi} E'$$

we get a correspondence: $\frac{E \xrightarrow[f]{\varphi} E'}{E \xrightarrow[id_B]{\tilde{\varphi}} p_*(f)(E')}$ such that for any polymaps $\psi : E_2 \rightarrow E_1$, $\xi : E'_1 \rightarrow E'_2$ and

$\varphi : E_1 \rightarrow E'_1$ we have $\xi \circ \varphi \circ \psi = p_*(f)(\xi) \circ \tilde{\varphi} \circ \psi$. In particular for $\psi = id_E$ and $\xi = id_{E'}$ then $\varphi = p_*(f)(id_{E'}) \circ \tilde{\varphi}$. This is the factorisation property of a weak cartesian morphism. A functor such that all the weak cartesian morphisms exists is a prefibration. It is a fibration if weak cartesian morphisms compose. This enforces for ∂p to be a pseudofunctor and not just lax.

Similarly asking for factorisation through $\mathbf{Cat}^{\text{op}} \hookrightarrow \mathbf{Dist}$ corresponds to representability of $\partial p(f)$ on its second variable while factorisation through $\mathbf{Adj} \hookrightarrow \mathbf{Dist}$ is representability in both variables. Combined with pseudofunctoriality this enforced ∂p to be an opfibration and a bifibration respectively.

Conversely, let $F : \mathcal{B} \rightarrow \mathbf{Dist}$ be a lax normal functor. We want to get a category $\int F$ and a functor $\int F \rightarrow \mathcal{B}$. This is often called the Grothendieck construction, especially when restricted to pseudofunctors $F : \mathcal{B} \rightarrow \mathbf{Cat}$. This can be obtained by considering the functor $\mathbf{Dist}_* \rightarrow \mathbf{Dist}$ from pointed distributors that

$$\begin{array}{ccc} \int F & \longrightarrow & \mathbf{Dist}_* \\ \text{forget about the point and consider the pullback} & \lrcorner & \downarrow \\ & & \mathcal{B} \longrightarrow \mathbf{Dist} \end{array} \quad \text{but for this to be well-defined there are}$$

some size considerations. Furthermore this will only give a lax functor $\int F \rightarrow \mathcal{B}$. That $\int F$ is a category has to be prove separately. We want to advertise another way to construct $\int F$ that is closer to the philosophy of this paper, using some notion of opcartesian 2-morphisms.

We will work in a slightly more general framework first. Let \mathcal{B} be a category, \mathcal{C} and \mathcal{D} be bicategories, $F : \mathcal{B} \rightarrow \mathcal{C}$ be a lax functor and $p : \mathcal{D} \rightarrow \mathcal{C}$ a strict functor. Furthermore suppose that for any 2-morphism $\alpha : f \Rightarrow g$ in \mathcal{C} and any morphism φ in \mathcal{D} lying over f , i.e. $p(\varphi) = f$, there is a morphism $\text{push}_\alpha(\varphi)$ over g and an opcartesian 2-morphism $\varphi \rightarrow \text{push}_\alpha(\varphi)$ lying over α . We now define a bicategory $\int_p F$. Its objects are pairs of objects (B, D) from \mathcal{B} and \mathcal{D} such that $F(B) = p(D)$. Similarly morphisms are pair of morphisms with the same image and 2-morphisms correspond to vertical 2-morphisms in \mathcal{D} . Given (B, D) an object of $\int_p F$ we want to define $id_{(B, D)} = (f, \varphi)$. We can take $f = id_B$. Then we have $F(f) = F(id_B)$. We would like to take $\varphi = id_D$ but $p(id_D) = id_{p(D)} = id_{F(B)} \neq F(id_B)$. However we have a 2-morphism $\eta_B : F(id_B) \Rightarrow id_{F(B)}$ from the lax functoriality of F . By using the fibrational property of p we can get an object over $id_{F(B)}$ by lifting id_D along η_B . So we have $id_{(B, D)} = (id_B, \text{push}_{\eta_B}(id_D))$. Similarly, $(g, \psi) \circ (f, \varphi) = (g \circ f, \text{push}_{\mu_{F(f), F(g)}}(\psi \circ \varphi))$ where $\mu_{F(f), F(g)} : F(g) \circ F(f) \Rightarrow F(g \circ f)$. Then we can prove that this are unital and associative up to isomorphism by using the same fact for \mathcal{B} and \mathcal{D} and the lax property of F and fibrational property of p . For example to prove right unitality of identity we consider:

$$\begin{array}{ccc}
 \begin{array}{c}
 D \xrightarrow{id_D} D \xrightarrow{\varphi} D' \\
 \downarrow \text{push}_\eta(id_D) \\
 \downarrow \text{push}_\mu(\varphi \circ \text{push}_\eta(id_D)) \\
 F(B) \xrightarrow{id_{F(B)}} F(B) \xrightarrow{F(f)} F(B') \\
 \downarrow \eta \\
 \downarrow \mu \\
 F(B) \xrightarrow{id_{F(B)}} F(B) \xrightarrow{F(f)} F(B') \\
 \downarrow F(id_B) \\
 \downarrow F(f)
 \end{array} & = & \begin{array}{c}
 D \xrightarrow{id_D} D \xrightarrow{\varphi} D' \\
 \downarrow unit_D \\
 \downarrow \varphi \\
 F(B) \xrightarrow{id_{F(B)}} F(B) \xrightarrow{F(f)} F(B') \\
 \downarrow unit_{F(B)} \\
 \downarrow F(f)
 \end{array}
 \end{array}$$

$unit_{F(B)}$ and $unit_D$ are 2-isomorphisms that comes from the bicategorical structure of \mathcal{C} and \mathcal{D} . The bottom equality comes from lax functoriality of F . Now using the universal property of cocartesian 2-morphisms we can factors $unit_D$ through $push_\eta(id_D)$ and then through the second pushforward to get a vertical 2-morphism $\chi : push_\mu(\varphi \circ push_\eta(id_D)) \xrightarrow{id_{F(f)}} \varphi$. And we get a vertical 2-morphism χ^{-1} in the other direction by precomposing

the diagram on the upper left corner with $unit_D^{-1}$. By definition we have $\chi \circ \chi^{-1} = unit_D \circ unit_D^{-1} = id_\varphi$. We get $\chi^{-1} \circ \chi = id_{push_\eta(\varphi \circ push_\mu(id_D))}$ by using the uniqueness of factorisation through a opcartesian 2-morphism. So left unitality of the identity holds up to vertical 2-isomorphism.

We get the following commutative diagram $\int_p F \longrightarrow \mathcal{D}$ by considering the projection functors from $\int_p F$.

$$\begin{array}{ccc}
 \int_p F & \longrightarrow & \mathcal{D} \\
 \downarrow & & \downarrow p \\
 \mathcal{B} & \xrightarrow{F} & \mathcal{C}
 \end{array}$$

We get back the Grothendieck construction by taking $p : \mathbf{Dist}_* \rightarrow \mathbf{Dist}$ the forgetful functor. Let show that it has opcartesian 2-morphism. Given $P, Q : A \rightarrow B$ and $\alpha : P \Rightarrow Q$ in \mathbf{Dist} , a morphism $(P, \varphi) : (A, a) \rightarrow (B, b)$ in \mathbf{Dist}_* over P is given by an element $\varphi \in P(a, b)$. Then we can use $\alpha_{a,b} : P(a, b) \rightarrow Q(a, b)$ to get a morphism $(Q, \alpha_{a,b}(\varphi)) : (A, a) \rightarrow (B, b)$ over Q with a 2-morphism $\alpha : (P, \varphi) \Rightarrow (Q, \alpha_{a,b}(\varphi))$. This is an opcartesian 2-morphism because for any 2-morphism in \mathbf{Dist} there is at most one 2-morphism over it for some given domain and codomain. So we get $\int F := \int_p F$ for this particular choice of p . If we expand the definition the objects of $\int F$ consists of pairs of objects (B, D) such that $B \in \mathcal{B}$, $D \in \mathbf{Dist}_*$ and $F(B) = p(D)$. Expanding a little more $D = (X, x)$ a pointed set such that $F(B) = X$. So it is just the data of a point in $F(B)$. Similarly, a morphism $(f, \varphi) : (B, x) \rightarrow (C, y)$ is a morphism $f : B \rightarrow C$ in \mathcal{B} and an element $\varphi \in F(f)(x, y)$. Finally a 2-morphism is a vertical 2-morphism in \mathbf{Dist}_* . But the only vertical morphisms are the identities. That makes $\int F$ a category.

These constructions are inverse to each other. First let start with $F : \mathcal{B} \rightarrow \mathbf{Dist}$. Let fix an object $B \in \mathcal{B}$. The fiber over B in $\int F$ is the set of objects (B, x) with $x \in F(B)$, which is isomorphic to the set $F(B)$. The profunctor $(\partial \int F)(f)$ is just $F(f)$. Conversely, given $p : \mathcal{E} \rightarrow \mathcal{B}$, $\int \partial p$ as for objects pairs (B, x) with $B \in \mathcal{B}$ and $x \in (\partial p)(B)$. But then $(\partial p)(B) = p^{-1}(B)$. So $Ob(\int \partial p) = \{(B, E) \mid p(E) = B\} = ob(\mathcal{E})$. Similarly $Hom_{\int \partial p}((B, E), (B', E')) = \{(f, \varphi) : B \times E \rightarrow B' \times E' \mid p(\varphi) = f\} = Hom_{\mathcal{E}}(E, E')$.